

DFTT 6/98
 NORDITA 98/10-HE
 KUL-TF-98/10
 February 1998

Microscopic string analysis of the D0-D8 brane system and dual R-R states*

M. Billó^{a†}, P. Di Vecchia^b, M. Frau^{c,f}, A. Lerda^{d,c,f},
 I. Pesando^{c,f}, R. Russo^{e,f} and S. Sciuto^{c,f}

^a *Instituut voor theoretische fysica,
 Katholieke Universiteit Leuven, B-3001 Leuven, Belgium*

^b *NORDITA, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark*

^c *Dipartimento di Fisica Teorica, Università di Torino*

^d *Dipartimento di Scienze e Tecnologie Avanzate
 Università di Torino, sede di Alessandria*

^e *Dipartimento di Fisica, Politecnico di Torino*

^f *I.N.F.N., Sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy*

Abstract

Using the boundary state formalism, we perform a microscopic string analysis of the interaction between two D-branes and provide a local interpretation for the R-R force in the D0-D8 brane system. To do so, we construct BRST invariant vertex operators for the massless R-R states in the asymmetric picture that are proportional to potentials rather than field strengths. The Hilbert space of such R-R states contains combinations of two vectors that decouple from all physical amplitudes, even in the presence of boundaries. Identifying these vectors, we remove the null states and recover duality relations among R-R potentials. If we specify to the D0-D8 brane system, this mechanism implies that the R-R 1-form state has a non-zero overlap with both the D0-brane and the D8-brane, thus explaining from a local point of view the non-vanishing R-R contribution in the interaction for the D0-D8 brane system and those related to it by duality.

*Work partially supported by the European Commission TMR programme ERBFMRX-CT96-0045 in which R.R. is associated to Torino University, and by MURST.

[†]e-mail: billo@tfdec1.fys.kuleuven.ac.be

1 Introduction

After the discovery [1] that D-branes are characterized by the fact that open strings with Dirichlet boundary conditions can end on them, it has been possible to study the interaction between two D-branes by computing a one-loop open string diagram with the open string stretching between the branes [1, 2]. An alternative way of describing D-branes is provided by the boundary state [3, 4, 5, 6, 7]. This is a BRST invariant state which can be interpreted as a source for a closed string emitted by a D-brane, and as such it is directly related to the classical brane solutions of the low-energy string effective action [8]. From this standpoint, the interaction between two D-branes is viewed as an exchange of closed string states, and thus it is computed with a tree-level diagram in which two boundary states are connected to each other by means of a closed string propagator. These two a priori completely independent approaches are actually equivalent and give exactly the same results, as a consequence of the modular properties of the string diagrams. From the explicit expression of the amplitude, one can easily see that when the two D-branes are near to each other, only the massless open string states are responsible for the interaction, whereas in the case of two distant D-branes only the massless closed string states give a non vanishing contribution.

If the two D-branes form a configuration which preserves enough space-time supersymmetry, then they do not interact. For example, this happens with two parallel Dp -branes, which break half of the supersymmetries. In this case, the vanishing of the force at the string level is a consequence of the “abstruse identity” and, from a field theory point of view, it can be understood [2] as due to a cancellation between the attractive contribution provided by the graviton and dilaton exchanges in the NS-NS sector, and the repulsive contribution provided by the exchange of a $(p+1)$ -form potential in the R-R sector. A zero force between two D-branes is also found in all other configurations in which the number ν of mixed Dirichlet-Neumann conditions for the open string stretching between the two D-branes is equal to 4 or 8 [1, 2, 9]. The case $\nu = 4$ has an explanation similar to the one we have just mentioned for parallel branes ($\nu = 0$). At the string level both the NS-NS and the R-R sectors are separately vanishing, and this is consistent with the fact that at large distance the graviton and the dilaton exactly compensate each other, while there is no massless R-R field which can be exchanged between the two D-branes since these couple to different R-R states.

On the contrary, the case $\nu = 8$ is more difficult to understand. At the string level, the vanishing of the force is again due to the “abstruse identity” which provides a cancellation between the non-zero contributions of the NS-NS and R-R sectors, but its microscopic interpretation is in this case problematic. In fact, the graviton and dilaton exchanges yield a repulsive force which should be compensated by an attractive one coming from the R-R potentials. However, from the field theory point of view, the branes of a $\nu = 8$ system seem to couple to different R-R states, and thus it appears impossible that they can exchange a R-R field. On the other

hand, it is undoubted that, when $\nu = 8$, there is a non-vanishing contribution from the massless R-R sector, which has also been used to explain the anomalous string creation when, for example, a D0-brane passes adiabatically through a D8-brane [7, 10, 11]. In fact, it turns out that after the two branes have exchanged their relative positions, the R-R force increases by a quantity proportional to the string tension $T = 1/(2\pi\alpha')$. This effect can then be interpreted as due to the creation of a fundamental string, and is related by a sequence of dualities to the creation of branes of different dimensions in other systems [12].

The puzzle of the field theory interpretation of the string results in the $\nu = 8$ system has been addressed and discussed in Ref. [10], where the unexpected R-R contribution is explained by treating the R-R 1-form emitted by a D0-brane as a background field for a D8-brane. Thus, the total R-R force can be directly read from the effective lagrangian as a non-local effect. An alternative explanation can be obtained by looking at the structure of the action of massive low-energy IIA supergravity [7, 13]. However, all these arguments are not on the same footing as the ones already discussed for the cases $\nu = 0, 4$, where the contributions of the various string sectors, that conspire to give the vanishing total result, are understood in terms of the exchanges of a dilaton, a graviton and a R-R state.

The main result of this paper is to show that also for the $\nu = 8$ systems it is possible to explain the R-R interaction from a microscopic and local point of view just like one usually does in all other cases. The crucial point is that the R-R charges of the two D-branes of a $\nu = 8$ system are essentially identified by a duality relation, and produce the same R-R potential. Therefore, the R-R interaction is nothing but the usual Coulomb-like force between D-branes.

More precisely in this paper we use the BRST invariant expression of the boundary state to compute, in the covariant formalism, the interaction between two D-branes as a closed string tree-level diagram. The contribution of the zero modes of the R-R sector turns out to be ill defined and hence, following Ref. [14], we introduce a regulator to obtain a meaningful result. In this way, we clearly see that the divergent superghost part and the vanishing matter contribution, when put together in the full amplitude, combine to leave a finite non-zero result if $\nu = 8$, while they give a vanishing result in all other cases. The final expression of the amplitude between two D-branes obtained in this way agrees with the one computed in Refs. [7, 9] without introducing the superghosts.

To give a microscopic interpretation of the force between two D-branes, it is necessary to know which closed string states can couple to the boundary state. While in the NS-NS sector there are no particular problems, it was already observed a few years ago in Ref. [15], that a boundary state has a non zero overlap only with R-R states in an asymmetric picture where the correspondent vertex operators contain the R-R potentials. On the contrary, the R-R states that are usually considered in perturbative string theory are in a symmetric picture and are proportional to the field strengths of the R-R potentials. To overcome this problem, we have explicitly constructed BRST invariant vertex operators for massless R-R states in the

asymmetric picture, and found that they are proportional to the R-R potentials, as suggested few years ago in Ref. [15], rather than to the corresponding field strengths. Furthermore, since the massless states exchanged between two D-branes cannot be on shell, it is necessary to study their off-shell properties. For the NS-NS sector it is known that the propagating states emitted by a D-brane are in the cohomology of a suitably restricted charge Q' [16]. We have found that for the R-R sector in the asymmetric picture there is a unique class of vertices W that are invariant under the restricted charge Q' also when extended off-shell. These new vertex operators contain an infinite number of terms, do not have left and right superghost number separately defined, and create states $|W\rangle$ that have exactly the same structure of the zero-mode part of the boundary states of the R-R sector. Despite the presence of infinite terms, these new states have a well-defined norm, provided that the scalar product is defined using the same regularization prescription introduced for boundary states. With this definition, a state $|W\rangle$ has a non vanishing scalar product not only with itself, but also with states $|W'\rangle$ carrying forms of different degree.

A careful analysis shows that there are two dimensional subspaces of the R-R Hilbert space with a degenerate metric. Thus, in each one of these subspaces, there exists a combination of two vectors which is null state, *i.e.* a state decoupling from all amplitudes, even with boundaries. This fact shows that in the Hilbert space of the asymmetric R-R sector there are pairs of dual vectors which describe the same state. On shell this is the usual Hodge duality between R-R potentials, whereas if we specify to the D0-D8 brane system, we see that, because of these identifications, the R-R 1-form state has a non-zero overlap with both the D0-brane and the D8-brane, thus solving the above-mentioned problem of the non vanishing R-R contribution to the interaction.

The paper is organized as follows. In Section 2, we give the full expression of the BRST invariant boundary state. Section 3 is devoted to the calculation of the static interaction between two D-branes. In Section 4 we construct the BRST invariant vertex operators in the asymmetric picture describing massless R-R potentials. In Section 5 we derive the duality relations between different R-R potentials and use them to show that for example a D0 and a D8 brane have opposite R-R charges. The factorization of the brane amplitude and its microscopic interpretation then follow immediately. Finally, in Appendix A we give some details on the construction of the BRST invariant asymmetric vertex operators for the R-R massless states with indefinite superghost number, and in Appendix B we show their BRST equivalence with vertex operators of zero left and right superghost number.

2 Boundary state for a Dp -brane

As explained in Ref. [8]¹, the boundary state $|B\rangle$ is a BRST invariant state of the closed string that inserts a boundary on the world-sheet and enforces the boundary

¹For an earlier discussion of the boundary state for the pure Neumann case see Refs. [14, 17].

conditions appropriate for a D-brane. For both the NS-NS and R-R sectors of the fermionic string, $|B\rangle$ can be written as the product of a matter part and a ghost part

$$|B\rangle = |B_{\text{mat}}\rangle |B_g\rangle \quad , \quad (2.1)$$

where

$$|B_{\text{mat}}\rangle = |B_X\rangle |B_\psi\rangle \quad , \quad |B_g\rangle = |B_{\text{gh}}\rangle |B_{\text{sgh}}\rangle \quad . \quad (2.2)$$

The matter part $|B_{\text{mat}}\rangle$ is defined by the overlap conditions that fix the identification at the boundary between the left and right movers of the matter fields X^μ and ψ^μ , namely

$$\partial_\tau X^\alpha|_{\tau=0}|B_X\rangle = 0 \quad , \quad (X^i - y^i)|_{\tau=0}|B_X\rangle = 0 \quad , \quad (2.3)$$

$$(\psi^\alpha - i\eta\tilde{\psi}^\alpha)|_{\tau=0}|B_\psi, \eta\rangle = 0 \quad , \quad (\psi^i + i\eta\tilde{\psi}^i)|_{\tau=0}|B_\psi, \eta\rangle = 0 \quad . \quad (2.4)$$

where α labels the $(p+1)$ Neumann (or longitudinal) directions, i labels the $(9-p)$ Dirichlet (or transverse) directions of a Dp -brane located at y . Notice that for ψ^μ there are two consistent identifications (corresponding to $\eta = \pm 1$), but, as we shall see, the GSO projection will allow only a superposition of the two.

By introducing the matrix

$$S_{\mu\nu} = (\eta_{\alpha\beta}, -\delta_{ij}) \quad , \quad (2.5)$$

and expanding the fields in modes, the overlap relations (2.3) and (2.4) become respectively

$$\begin{aligned} (\alpha_n + S \cdot \tilde{\alpha}_{-n})|B_X\rangle &= 0 \quad (n \neq 0) \quad , \\ \hat{p}^\alpha|B_X\rangle &= (\hat{q}^i - y^i)|B_X\rangle = 0 \quad , \end{aligned} \quad (2.6)$$

and

$$(\psi_m - i\eta S \cdot \tilde{\psi}_{-m})|B_\psi, \eta\rangle = 0 \quad (2.7)$$

where the index m is integer in the R sector and half-integer in the NS sector.

It is not difficult to check that the identifications (2.6) and (2.7) imply that $|B_{\text{mat}}, \eta\rangle$ is annihilated by the following linear combinations of left and right generators of the super Virasoro algebra

$$(L_n^{\text{mat}} - \tilde{L}_{-n}^{\text{mat}})|B_{\text{mat}}, \eta\rangle = 0 \quad , \quad (G_m^{\text{mat}} + i\eta\tilde{G}_{-m}^{\text{mat}})|B_{\text{mat}}, \eta\rangle = 0 \quad . \quad (2.8)$$

Since the boundary state $|B, \eta\rangle$ must be BRST invariant, that is²

$$(Q + \tilde{Q})|B, \eta\rangle = 0 \quad , \quad (2.9)$$

²The boundary state must be in the cohomology of the total BRST charge $Q + \tilde{Q}$; contrarily to the usual perturbative states of the closed string, it cannot be chosen to be annihilated separately by Q and $\tilde{Q}$ due to the presence of a boundary on the world-sheet.

the relations (2.8) must be supplemented by the analogous ones in the ghost sector, namely

$$\left(L_n^g - \tilde{L}_{-n}^g\right) |B_g, \eta\rangle = 0 \quad , \quad \left(G_m^g + i\eta \tilde{G}_{-m}^g\right) |B_g, \eta\rangle = 0 \quad . \quad (2.10)$$

They imply that

$$\begin{aligned} (c_n + \tilde{c}_{-n}) |B_{gh}\rangle &= 0 \quad , \quad (b_n - \tilde{b}_{-n}) |B_{gh}\rangle = 0 \quad , \\ (\gamma_m + i\eta \tilde{\gamma}_{-m}) |B_{sgh}, \eta\rangle &= 0 \quad , \quad (\beta_m + i\eta \tilde{\beta}_{-m}) |B_{sgh}, \eta\rangle = 0 \quad . \end{aligned} \quad (2.11)$$

Using the conventions and normalizations of Ref. [8], we can write the solution to the overlap equations (2.6), (2.7) and (2.11) as follows

$$|B, \eta\rangle_{R, NS} = \frac{T_p}{2} |B_X\rangle |B_{gh}\rangle |B_\psi, \eta\rangle_{R, NS} |B_{sgh}, \eta\rangle_{R, NS} \quad , \quad (2.12)$$

where

$$|B_X\rangle = \delta^{(d_\perp)}(\hat{q} - y) \exp\left[-\sum_{n=1}^{\infty} \frac{1}{n} \alpha_{-n} \cdot S \cdot \tilde{\alpha}_{-n}\right] |0; k=0\rangle \quad , \quad (2.13)$$

$$|B_{gh}\rangle = \exp\left[\sum_{n=1}^{\infty} (c_{-n} \tilde{b}_{-n} - b_{-n} \tilde{c}_{-n})\right] \frac{c_0 + \tilde{c}_0}{2} |q=1\rangle |\tilde{q}=1\rangle \quad , \quad (2.14)$$

and, in the NS sector in the $(-1, -1)$ picture,

$$|B_\psi, \eta\rangle_{NS} = \exp\left[i\eta \sum_{m=1/2}^{\infty} \psi_{-m} \cdot S \cdot \tilde{\psi}_{-m}\right] |0\rangle \quad , \quad (2.15)$$

$$|B_{sgh}, \eta\rangle_{NS} = \exp\left[i\eta \sum_{m=1/2}^{\infty} (\gamma_{-m} \tilde{\beta}_{-m} - \beta_{-m} \tilde{\gamma}_{-m})\right] |P=-1\rangle |\tilde{P}=-1\rangle \quad , \quad (2.16)$$

or, in the R sector in the $(-1/2, -3/2)$ picture,

$$|B_\psi, \eta\rangle_R = \exp\left[i\eta \sum_{m=1}^{\infty} \psi_{-m} \cdot S \cdot \tilde{\psi}_{-m}\right] |B_\psi, \eta\rangle_R^{(0)} \quad , \quad (2.17)$$

$$|B_{sgh}, \eta\rangle_R = \exp\left[i\eta \sum_{m=1}^{\infty} (\gamma_{-m} \tilde{\beta}_{-m} - \beta_{-m} \tilde{\gamma}_{-m})\right] |B_{sgh}, \eta\rangle_R^{(0)} \quad , \quad (2.18)$$

where the superscript $^{(0)}$ denotes the zero-mode contribution to be discussed momentarily. The overall normalization factor T_p can be unambiguously fixed from the factorization of amplitudes of closed strings emitted from a disk [5, 8] and is the tension of the Dp -brane

$$T_p = \sqrt{\pi} (2\pi \sqrt{\alpha'})^{3-p} \quad . \quad (2.19)$$

To write explicitly the zero-mode parts of the boundary state in the R-R sector, it is necessary to introduce some further notation. Let

$$|A\rangle |\tilde{B}\rangle = \lim_{z, \bar{z} \rightarrow 0} S^A(z) \tilde{S}^B(\bar{z}) |0\rangle \quad (2.20)$$

denote the vacuum for the fermionic zero-modes ψ_0^μ and $\tilde{\psi}_0^\mu$, where S^A and $\tilde{S}^B$ are the spin fields in the 32-dimensional Majorana representation; then, as shown in Ref. [8], we have

$$|B_\psi, \eta\rangle_{\text{R}}^{(0)} = \mathcal{M}_{AB}^{(\eta)} |A\rangle |\tilde{B}\rangle \quad , \quad (2.21)$$

where

$$\mathcal{M}^{(\eta)} = C \Gamma^0 \Gamma^{l_1} \dots \Gamma^{l_p} \left(\frac{1 + i\eta \Gamma_{11}}{1 + i\eta} \right) \quad , \quad (2.22)$$

with C being the charge conjugation matrix and l_i labeling the space directions of the D-brane world volume. Finally, if $|P = -1/2\rangle |\tilde{P} = -3/2\rangle$ denotes the superghost vacuum in the $(-1/2, -3/2)$ picture that is annihilated by β_0 and $\tilde{\gamma}_0$, we have [14]

$$|B_{\text{sgh}}, \eta\rangle_{\text{R}}^{(0)} = \exp[i\eta \gamma_0 \tilde{\beta}_0] |P = -1/2\rangle |\tilde{P} = -3/2\rangle \quad . \quad (2.23)$$

Before using the boundary state to compute amplitudes involving D-branes, one must perform the GSO projection. In the NS-NS sector the projected state is

$$|B\rangle_{\text{NS}} \equiv \frac{1 - (-1)^{F+G}}{2} \frac{1 - (-1)^{\tilde{F}+\tilde{G}}}{2} |B, +\rangle_{\text{NS}} \quad , \quad (2.24)$$

where F and G are the fermion and superghost number operators

$$F = \sum_{m=1/2}^{\infty} \psi_{-m} \cdot \psi_m \quad , \quad G = - \sum_{m=1/2}^{\infty} (\gamma_{-m} \beta_m + \beta_{-m} \gamma_m) \quad . \quad (2.25)$$

After some simple algebra, it is easy to see that

$$|B\rangle_{\text{NS}} = \frac{1}{2} (|B, +\rangle_{\text{NS}} - |B, -\rangle_{\text{NS}}) \quad (2.26)$$

In the R-R sector the GSO projected boundary state is

$$|B\rangle_{\text{R}} \equiv \frac{1 + (-1)^p (-1)^{F+G}}{2} \frac{1 - (-1)^{\tilde{F}+\tilde{G}}}{2} |B, +\rangle_{\text{R}} \quad . \quad (2.27)$$

where p is even for Type IIA and odd for Type IIB, and

$$(-1)^F = \Gamma_{11} (-1)^{\sum_{m=1}^{\infty} \psi_{-m} \cdot \psi_m} \quad , \quad G = -\gamma_0 \beta_0 - \sum_{m=1}^{\infty} [\gamma_{-m} \beta_m + \beta_{-m} \gamma_m] \quad . \quad (2.28)$$

After some straightforward manipulations, one can check that

$$|B\rangle_{\text{R}} = \frac{1}{2} (|B, +\rangle_{\text{R}} + |B, -\rangle_{\text{R}}) \quad . \quad (2.29)$$

For later convenience we now rewrite the boundary state $|B\rangle_{\text{R}}$ using 16-dimensional chiral and antichiral spinor indices α and $\dot{\alpha}$ for Majorana-Weyl fermions. Then, for the Type IIA theory we have

$$\begin{aligned} |B\rangle_{\text{R}} = & \frac{T_p}{2} |B_X\rangle |B_{\text{gh}}\rangle \left\{ \left(C\Gamma^0\Gamma^{l_1} \dots \Gamma^{l_p} \right)_{\alpha\beta} \cos [\gamma_0\tilde{\beta}_0 + \Theta] |\alpha\rangle_{-1/2} |\tilde{\beta}\rangle_{-3/2} \right. \\ & \left. + \left(C\Gamma^0\Gamma^{l_1} \dots \Gamma^{l_p} \right)_{\dot{\alpha}\dot{\beta}} \sin [\gamma_0\tilde{\beta}_0 + \Theta] |\dot{\alpha}\rangle_{-1/2} |\tilde{\dot{\beta}}\rangle_{-3/2} \right\} , \end{aligned} \quad (2.30)$$

where

$$\Theta = \sum_{m=1}^{\infty} (\psi_{-m} \cdot S \cdot \tilde{\psi}_{-m} + \gamma_{-m} \tilde{\beta}_{-m} - \beta_{-m} \tilde{\gamma}_{-m}) , \quad (2.31)$$

and we have abbreviated $|\alpha, P = \ell\rangle$ with $|\alpha\rangle_{\ell}$. Note that one also has the following identity

$$|\alpha\rangle_{\ell} = \lim_{z \rightarrow 0} S^{\alpha}(z) e^{\ell\phi(z)} |0\rangle . \quad (2.32)$$

where ϕ is the chiral boson of the superghost fermionization formulas

$$\gamma(z) = e^{\phi(z)} \eta(z) , \quad \beta(z) = \partial\xi(z) e^{-\phi(z)} . \quad (2.33)$$

The boundary state for the Type IIB theory can be simply obtained from Eq. (2.30) by changing the chirality of the left moving fermions according to Eq. (2.27), and explicitly reads

$$\begin{aligned} |B\rangle_{\text{R}} = & \frac{T_p}{2} |B_X\rangle |B_{\text{gh}}\rangle \left\{ \left(C\Gamma^0\Gamma^{l_1} \dots \Gamma^{l_p} \right)_{\dot{\alpha}\dot{\beta}} \cos [\gamma_0\tilde{\beta}_0 + \Theta] |\dot{\alpha}\rangle_{-1/2} |\tilde{\dot{\beta}}\rangle_{-3/2} \right. \\ & \left. + \left(C\Gamma^0\Gamma^{l_1} \dots \Gamma^{l_p} \right)_{\alpha\beta} \sin [\gamma_0\tilde{\beta}_0 + \Theta] |\alpha\rangle_{-1/2} |\tilde{\beta}\rangle_{-3/2} \right\} , \end{aligned} \quad (2.34)$$

with p odd.

We would like to stress that the boundary states $|B\rangle_{\text{NS,R}}$ are written in a definite picture $(P, \tilde{P})$ of the superghost system, where

$$P = \oint \frac{dz}{2\pi i} (-\partial\phi + \xi\eta) \quad (2.35)$$

and $\tilde{P} = -2 - P$ in order to soak up the anomaly in the superghost number. In particular we have chosen $P = -1$ in the NS sector and $P = -1/2$ in the R sector, even if other choices would have been possible in principle [14]. Since P is half-integer in the R sector, the boundary state $|B\rangle_{\text{R}}$ has always $P \neq \tilde{P}$, and thus, as suggested in Ref. [15], it can couple only to R-R states in the asymmetric picture $(P, \tilde{P})$. However, the crucial point is to observe that the massless R-R states in the $(-1/2, -3/2)$ picture may contain a part that is proportional to the R-R *potentials*, as opposed to the standard massless R-R states in the symmetric picture $(-1/2, -1/2)$ that are always proportional to the R-R field strengths. This property, which has been already exploited in Ref. [8] to derive the D-brane effective

action and the long-distance behavior of the D-brane solutions from the boundary state formalism, will be used later on to study the interaction between two D-branes. Many details are contained in Section 4 and in the Appendices. Finally, we observe that it is possible to fermionize the boundary states $|B\rangle_{\text{NS,R}}$ using Eq. (2.33). However, as shown in Ref. [14] the equivalent boundary states in the (ϕ, η, ξ) system contain an infinite number of terms corresponding to the infinite possibilities of satisfying (2.35).

We conclude this section by writing the conjugate boundary state

$${}_{\text{R,NS}}\langle B, \eta| = \frac{T_p}{2} \langle B_X| \langle B_{\text{gh}}| {}_{\text{R,NS}}\langle B_\psi, \eta| {}_{\text{R,NS}}\langle B_{\text{sgh}}, \eta| \quad (2.36)$$

which we will use in the calculation of the amplitudes between D-branes. The explicit expressions for the various factors in Eq. (2.36) can be obtained by solving the overlap relations for the conjugate boundary state and are given by the obvious counterparts of Eqs. (2.13)-(2.18). For example the fermion and the superghost contributions are

$${}_{\text{NS}}\langle B_\psi, \eta| = \langle 0| \exp\left[-i\eta \sum_{m=1/2}^{\infty} \psi_m \cdot S \cdot \tilde{\psi}_m\right] \quad (2.37)$$

and

$${}_{\text{NS}}\langle B_{\text{sgh}}, \eta| = \langle P = -1| \langle \tilde{P} = -1| \exp\left[i\eta \sum_{m=1/2}^{\infty} (\beta_m \tilde{\gamma}_m - \gamma_m \tilde{\beta}_m)\right] \quad (2.38)$$

in the NS-NS sector, and

$${}_{\text{R}}\langle B_\psi, \eta| = \langle A| \langle \tilde{B}| \mathcal{N}_{AB}^{(\eta)} \exp\left[-i\eta \sum_{m=1}^{\infty} \psi_m \cdot S \cdot \tilde{\psi}_m\right] , \quad (2.39)$$

with

$$\mathcal{N}^{(\eta)} = \Gamma^{0T} \mathcal{M}^{(\eta)} \Gamma^0 = (-1)^p C \Gamma^0 \Gamma^{l_1} \dots \Gamma^{l_p} \left(\frac{1 - i\eta \Gamma_{11}}{1 + i\eta} \right) , \quad (2.40)$$

and

$${}_{\text{R}}\langle B_{\text{sgh}}, \eta| = \langle P = -3/2| \langle \tilde{P} = -1/2| \exp[i\eta \beta_0 \tilde{\gamma}_0] \exp\left[i\eta \sum_{m=1}^{\infty} (\beta_m \tilde{\gamma}_m - \gamma_m \tilde{\beta}_m)\right] \quad (2.41)$$

in the R-R sector.

3 Interaction between a p and a p' brane

In this section we study the static interaction between a D-brane located at y_1 , and a D-brane located at y_2 , with NN directions common to the brane world-volumes,

$DD \geq 1$ directions transverse to both, and $\nu = (10 - NN - DD)$ directions of mixed type. We will not consider instantonic D-branes [18], hence also $NN \geq 1$. The two D-branes simply interact via tree-level exchange of closed strings whose propagator is

$$D = \frac{\alpha'}{4\pi} \int \frac{d^2 z}{|z|^2} z^{L_0} \bar{z}^{\tilde{L}_0} , \quad (3.1)$$

so that the static amplitude is given by

$$\mathcal{A} = \langle B^1 | D | B^2 \rangle , \quad (3.2)$$

where $|B^1\rangle$ and $|B^2\rangle$ are the boundary states describing the two D-branes³.

The evaluation of $\mathcal{A}$ in the NS-NS sector presents no difficulties and can be performed starting from the definitions given in Section 2 and using standard string techniques. Here we simply quote the final result, namely

$$\begin{aligned} \mathcal{A}_{\text{NS-NS}} &= \frac{V_{NN}}{2\pi} (8\pi^2 \alpha')^{-\frac{NN}{2}} \int_0^\infty dt \left(\frac{\pi}{t} \right)^{\frac{DD}{2}} e^{-\Delta Y^2 / (2\alpha' t)} \\ &\times \left[\left(\frac{f_3}{f_1} \right)^{8-\nu} \left(\frac{f_4}{f_2} \right)^\nu - \left(\frac{f_4}{f_1} \right)^{8-\nu} \left(\frac{f_3}{f_2} \right)^\nu \right] , \end{aligned} \quad (3.3)$$

where V_{NN} is the common world-volume of the two D-branes, ΔY is the transverse distance between them, and the functions f_i are, as usual, given by

$$\begin{aligned} f_1 &= q^{\frac{1}{12}} \prod_{n=1}^\infty (1 - q^{2n}) , & f_2 &= \sqrt{2} q^{\frac{1}{12}} \prod_{n=1}^\infty (1 + q^{2n}) , \\ f_3 &= q^{-\frac{1}{24}} \prod_{n=1}^\infty (1 + q^{2n-1}) , & f_4 &= q^{-\frac{1}{24}} \prod_{n=1}^\infty (1 - q^{2n-1}) , \end{aligned} \quad (3.4)$$

with $q = e^{-t}$. It is interesting to notice that the two terms in the square brackets of Eq. (3.3) come respectively from the NS-NS $(-1)^{(F+G)}$ and the NS-NS sectors of the exchanged closed string, which, under the transformation $t \rightarrow 1/t$, are mapped into the NS and R sectors of the open string suspended between the branes. Notice that $\mathcal{A}_{\text{NS-NS}} = 0$ if $\nu = 4$.

On the contrary, the evaluation of $\mathcal{A}$ in the R-R sector requires more care due to the presence of zero-modes in both the fermionic matter fields and the bosonic superghosts. To perform the calculation it is convenient to use boundary states before the GSO projection, and thus consider

$$\mathcal{A}_{\text{R-R}}(\eta_1, \eta_2) = {}_{\text{R}}\langle B^1, \eta_1 | D | B^2, \eta_2 \rangle_{\text{R}} . \quad (3.5)$$

³In Eq. (3.2) it is understood the usual insertion of the (b, c) zero-modes that is necessary to soak up the ghost anomaly of the cylinder diagram (see for example Ref. [17]).

As before, it is not difficult to compute the contribution of X^μ , (b, c) and also of the non-zero-modes of ψ^μ and (β, γ) . Indeed, after some algebra, we find

$$\begin{aligned} \mathcal{A}_{\text{R-R}}(\eta_1, \eta_2) &= \frac{V_{NN}}{\pi} (8\pi^2 \alpha')^{-\frac{NN}{2}} 2^{-\frac{\nu}{2}} \int_0^\infty dt \left(\frac{\pi}{t} \right)^{\frac{DD}{2}} e^{-\Delta Y^2 / (2\alpha' t)} \\ &\times \left[2^{\nu-4} \left(\frac{f_2}{f_1} \right)^{8-2\nu} \delta_{\eta_1 \eta_2, -1} + \delta_{\eta_1 \eta_2, +1} \right] {}^{(0)}_{\text{R}} \langle B^1, \eta_1 | B^2, \eta_2 \rangle_{\text{R}}^{(0)}, \end{aligned} \quad (3.6)$$

where

$$|B, \eta\rangle_{\text{R}}^{(0)} = |B_\psi, \eta\rangle_{\text{R}}^{(0)} |B_{\text{sgh}}, \eta\rangle_{\text{R}}^{(0)} \quad (3.7)$$

(see Eqs. (2.21) and (2.23)). Note that in Eq. (3.6) it is essential *not* to separate the matter and the superghost zero-modes. In fact, a naïve evaluation of ${}^{(0)}_{\text{R}} \langle B^1, \eta_1 | B^2, \eta_2 \rangle_{\text{R}}^{(0)}$ would lead to a divergent or ill defined result: after expanding the exponentials in ${}^{(0)}_{\text{R}} \langle B_{\text{sgh}}^1, \eta_1 | B_{\text{sgh}}^2, \eta_2 \rangle_{\text{R}}^{(0)}$, all the infinite terms with any superghost number contribute, and yield the divergent sum $1 + 1 + 1 + \dots$ if $\eta_1 \eta_2 = 1$, or the alternating sum $1 - 1 + 1 - \dots$ if $\eta_1 \eta_2 = -1$. This problem has already been addressed in Ref. [14] and solved by introducing a regularization scheme for the pure Neumann case ($NN = 10$). Here, we propose the extension of this method to the most general case with D-branes.

We define the scalar product in Eq. (3.6) as follows

$${}^{(0)}_{\text{R}} \langle B^1, \eta_1 | B^2, \eta_2 \rangle_{\text{R}}^{(0)} \equiv \lim_{x \rightarrow 1} {}^{(0)}_{\text{R}} \langle B^1, \eta_1 | \mathcal{R}(x) | B^2, \eta_2 \rangle_{\text{R}}^{(0)}, \quad (3.8)$$

where the regulator is

$$\mathcal{R}(x) = x^{2(F_0 + G_0)} \quad (3.9)$$

where F_0 and G_0 are the zero-mode parts of the operators F and G (implicitly) defined in Eq. (2.28). After inserting the regulator, we can factorize the scalar product and write

$$\begin{aligned} {}^{(0)}_{\text{R}} \langle B^1, \eta_1 | B^2, \eta_2 \rangle_{\text{R}}^{(0)} &= \lim_{x \rightarrow 1} \left[{}^{(0)}_{\text{R}} \langle B_\psi^1, \eta_1 | x^{2F_0} | B_\psi^2, \eta_2 \rangle_{\text{R}}^{(0)} \right. \\ &\times \left. {}^{(0)}_{\text{R}} \langle B_{\text{sgh}}^1, \eta_1 | x^{2G_0} | B_{\text{sgh}}^2, \eta_2 \rangle_{\text{R}}^{(0)} \right]. \end{aligned} \quad (3.10)$$

Let us first consider the superghost part. Remembering that $G_0 = -\gamma_0 \beta_0$, we simply have

$$\begin{aligned} {}^{(0)}_{\text{R}} \langle B_{\text{sgh}}^1, \eta_1 | x^{2G_0} | B_{\text{sgh}}^2, \eta_2 \rangle_{\text{R}}^{(0)} &= \langle -3/2, -1/2 | e^{i\eta_1 \beta_0 \tilde{\gamma}_0} x^{-2\gamma_0 \beta_0} e^{i\eta_2 \gamma_0 \tilde{\beta}_0} | -1/2, -3/2 \rangle \\ &= \frac{1}{1 - \eta_1 \eta_2 x^2}. \end{aligned} \quad (3.11)$$

To discuss the regularization of the fermionic part, it is convenient (even though not necessary) to perform a Wick rotation $\Gamma^0 \rightarrow i\Gamma^{10}$ and work with the Γ matrices

of $SO(10)$. Then, we can group them into five pairs, and for each pair $\mathbf{a} = (a_1, a_2)$, introduce fermionic annihilation and creation operators

$$e_{\mathbf{a}}^{\pm} = \frac{\Gamma^{a_1} \pm i\Gamma^{a_2}}{2} , \quad (3.12)$$

satisfying

$$\{e_{\mathbf{a}}^+, e_{\mathbf{b}}^-\} = \delta_{\mathbf{a}, \mathbf{b}} . \quad (3.13)$$

The Hilbert space associated to each couple of Γ matrices is two-dimensional, and it is spanned by the states $|\uparrow\rangle_{\mathbf{a}}$ and $|\downarrow\rangle_{\mathbf{a}} = e_{\mathbf{a}}^- |\uparrow\rangle_{\mathbf{a}}$, which are eigenvectors of the number operator

$$N_{\mathbf{a}} = [e_{\mathbf{a}}^+, e_{\mathbf{a}}^-] = -i\Gamma^{a_1}\Gamma^{a_2} \quad (3.14)$$

with eigenvalues $+1$ and -1 respectively. Thus, the 32-dimensional space spanned by a $SO(10)$ Majorana spinor is expressed as the direct product of five copies of these two-dimensional spaces, each copy for two different directions. Note that in this way we explicitly break $SO(10) \rightarrow SO(2)^5$, but if all directions are equivalent, the final results will still be $SO(10)$ invariant (see Ref. [14]). In the case we are considering, not all directions are on the same footing because of the different boundary conditions imposed by the D-branes. Therefore, it is necessary to exert some care in pairing the space directions and the corresponding Γ matrices if we want to have meaningful final results. To this aim it is necessary that directions of mixed type be grouped together, while the remaining NN or DD directions can be paired as one wishes. However, since for all configurations we are considering, the time direction (*i.e.* the 10th after Wick rotation) is always NN, and the 9th direction can always be chosen as DD, we group them together, so that the prescription can always be used. Note that this choice amounts to specify the space direction in which the light-cone is oriented (namely the 9th), or equivalently to mark in the covariant formalism the direction which together with the time is “canceled” by the superghosts.

For the sake of simplicity, we now suppose that the NN directions of the two D-branes are $(0, \dots, p')$ and $(0, \dots, p)$ with $p' \leq p$. This specific choice allows us to perform explicitly the calculation, but it is not restrictive since any other configuration of D-branes can be reduced to it by a sequence of T-duality and parity transformations. Then, we group the ten Γ matrices in five pairs, for example as follows

$$\begin{aligned} \text{IIA} &\longrightarrow (1\,2)(3\,4)(5\,6)(7\,8)(9\,10) , \\ \text{IIB} &\longrightarrow (2\,3)(4\,5)(6\,7)(1\,8)(9\,10) . \end{aligned} \quad (3.15)$$

The next task is to find the expression for F_0 in this basis. To do so, we observe that, in the ten dimensional euclidean space [14],

$$(-1)^{F_0} = \prod_{\mu=1}^{10} \Gamma^{\mu} = i \prod_{k=1}^5 N_k , \quad (3.16)$$

where $N_k \equiv N_{\mathbf{a}_k}$. Then, since $\exp(i N_k \pi/2) = i N_k$, we can rewrite Eq. (3.16) as follows

$$(-1)^{F_0} = \prod_{k=1}^5 \exp\left(i N_k \frac{\pi}{2}\right) , \quad (3.17)$$

from which we can define

$$F_0 = \frac{1}{2} \sum_{k=1}^5 N_k . \quad (3.18)$$

Thus, the regulator for the fermionic zero-modes is

$$x^{2F_0} = x^{\sum_{k=1}^5 N_k} . \quad (3.19)$$

We are now in the position of computing explicitly the contribution of the fermionic zero-modes to the D-brane amplitude. Indeed, we have

$${}^{(0)}_{\text{R}} \langle B_{\psi}^1, \eta_1 | x^{2F_0} | B_{\psi}^2, \eta_2 \rangle_{\text{R}}^{(0)} = \text{tr} \left(x^{2F_0} \mathcal{M}^{(\eta_2)} C^{-1} \mathcal{N}^{(\eta_1)T} C^{-1} \right) , \quad (3.20)$$

where, to obtain the right hand side, we have used the inner product

$$\left(\langle A | \langle \tilde{B} | \right) \left(| D \rangle | \tilde{E} \rangle \right) = - \langle A | D \rangle \langle \tilde{B} | \tilde{E} \rangle = -(C^{-1})^{AD} (C^{-1})^{BE} \quad (3.21)$$

with the minus sign due to the exchange in the ordering of the spinor states. Inserting the explicit definitions (2.22) and (2.40) of the matrices $\mathcal{M}$ and $\mathcal{N}$, after some straightforward algebra, we find

$${}^{(0)}_{\text{R}} \langle B_{\psi}^1, \eta_1 | x^{2F_0} | B_{\psi}^2, \eta_2 \rangle_{\text{R}}^{(0)} = - \text{tr} \left(x^{2F_0} \prod_{\alpha} \Gamma^{\alpha} \right) \delta_{\eta_1 \eta_2, -1} - (-1)^p \text{tr} \left(x^{2F_0} \prod_{\alpha} \Gamma^{\alpha} \Gamma_{11} \right) \delta_{\eta_1 \eta_2, +1} \quad (3.22)$$

where the index α runs over the ν directions of mixed type. By introducing the fermionic number operators as in Eq. (3.14) and recalling that $\Gamma_{11} = \prod_k N_k$, we can easily compute the traces and get

$$\begin{aligned} \text{tr} \left(x^{2F_0} \prod_{\alpha} \Gamma^{\alpha} \right) &= \prod_{k=1}^{\frac{10-\nu}{2}} \text{tr} \left(x^{N_k} \right) \prod_{l=1}^{\frac{\nu}{2}} \text{tr} \left(i x^{N_l} N_l \right) \\ &= i^{\frac{\nu}{2}} \left(x + \frac{1}{x} \right)^{\frac{10-\nu}{2}} \left(x - \frac{1}{x} \right)^{\frac{\nu}{2}} , \end{aligned} \quad (3.23)$$

$$\begin{aligned} \text{tr} \left(x^{2F_0} \prod_{\alpha} \Gamma^{\alpha} \Gamma_{11} \right) &= \prod_{l=1}^{\frac{\nu}{2}} \text{tr} \left(i x^{N_l} \right) \prod_{k=1}^{\frac{10-\nu}{2}} \text{tr} \left(x^{N_k} N_k \right) \\ &= i^{\frac{\nu}{2}} \left(x + \frac{1}{x} \right)^{\frac{\nu}{2}} \left(x - \frac{1}{x} \right)^{\frac{10-\nu}{2}} . \end{aligned} \quad (3.24)$$

Putting together the fermionic and superghost contributions as in Eq. (3.10), we finally obtain

$$\begin{aligned}
{}_{\text{R}}^{(0)}\langle B^1, \eta_1 | B^2, \eta_2 \rangle_{\text{R}}^{(0)} &= \lim_{x \rightarrow 1} \left[-i^{\frac{\nu}{2}} \left(x + \frac{1}{x} \right)^{\frac{10-\nu}{2}} \left(x - \frac{1}{x} \right)^{\frac{\nu}{2}} \frac{1}{1+x^2} \delta_{\eta_1 \eta_2, -1} \right. \\
&\quad \left. - (-1)^p i^{\frac{\nu}{2}} \left(x + \frac{1}{x} \right)^{\frac{\nu}{2}} \left(x - \frac{1}{x} \right)^{\frac{10-\nu}{2}} \frac{1}{1-x^2} \delta_{\eta_1 \eta_2, +1} \right] \\
&= -16 \delta_{\nu, 0} \delta_{\eta_1 \eta_2, -1} + 16 \delta_{\nu, 8} \delta_{\eta_1 \eta_2, +1} .
\end{aligned} \tag{3.25}$$

In our configuration, $\nu = 8$ can be realized only for $p' = 0$ and $p = 8$ so that we can understand the factor of $(-1)^p$; note that in this case there is a cancellation between the zero coming from the fermionic zero-modes and the infinity coming from the superghost zero-modes. This cancellation was qualitatively suggested in the Appendix of Ref. [7], here we give a more rigorous derivation of it. It is interesting to observe that Eq. (3.25) implies that the R-R part of the D-brane amplitude vanishes unless the two branes are parallel ($\nu = 0$) or maximally orthogonal ($\nu = 8$) to each other. Furthermore, the zero-mode contribution to the R-R amplitude, which is the only one that survives in the field theory limit, is numerically equal in both cases, and in particular it is repulsive for $\nu = 0$ and attractive for $\nu = 8$. This fact suggests that the nature of the force is the same in the two cases and that the charges of two D-branes with $\nu = 8$ can be essentially identified; in Section 5 we will show that this interpretation is indeed correct.

We now write the final expression for the R-R amplitude. Inserting Eq. (3.25) into Eq. (3.6), we get

$$\begin{aligned}
\mathcal{A}_{\text{R-R}}(\eta_1, \eta_2) &= \frac{V_{NN}}{\pi} (8\pi^2 \alpha')^{-\frac{NN}{2}} \int_0^\infty dt \left(\frac{\pi}{t} \right)^{\frac{DD}{2}} e^{-\Delta Y^2 / (2\alpha' t)} \\
&\quad \times \left[- \left(\frac{f_2}{f_1} \right)^8 \delta_{\nu, 0} \delta_{\eta_1 \eta_2, -1} + \delta_{\nu, 8} \delta_{\eta_1 \eta_2, +1} \right] ,
\end{aligned} \tag{3.26}$$

from which we immediately obtain the GSO projected amplitude

$$\mathcal{A}_{\text{R-R}} = \frac{V_{NN}}{2\pi} (8\pi^2 \alpha')^{-\frac{NN}{2}} \int_0^\infty dt \left(\frac{\pi}{t} \right)^{\frac{DD}{2}} e^{-\Delta Y^2 / (2\alpha' t)} \left[- \left(\frac{f_2}{f_1} \right)^8 \delta_{\nu, 0} + \delta_{\nu, 8} \right] . \tag{3.27}$$

We conclude this section with a few observations. First of all, the final result $\mathcal{A}_{\text{R-R}}$ is Lorentz invariant, even if we used a non Lorentz invariant regularization prescription to compute the zero-mode contribution. Secondly, the $\nu = 0$ and $\nu = 8$ terms in Eq. (3.27) come respectively from the R-R $(-1)^{(F+G)}$ and the R-R sectors of the exchanged closed string, which, under the transformation $t \rightarrow 1/t$, are mapped into the NS $(-1)^{(F+G)}$ and R $(-1)^{(F+G)}$ sectors of the open string suspended between the branes. Moreover, due to the “abstruse identity”, the total D-brane amplitude

$$\mathcal{A} = \mathcal{A}_{\text{NS-NS}} + \mathcal{A}_{\text{R-R}} \tag{3.28}$$

vanishes if $\nu = 0, 4, 8$; these are precisely the configurations of two D-branes which break half of the supersymmetries of the Type II theory and satisfy the BPS no-force condition.

Note that for the $\nu = 8$ systems the repulsive NS-NS force is canceled by the attractive contribution of the R-R sector that contains an insertion of Γ_{11} (see Eqs. (3.22) and (3.24)). Since it is not parity invariant, this contribution becomes repulsive if a parity transformation along the DD direction (*i.e.* X^9) is performed. As pointed out in Ref. [19], such a parity transformation maps the original $\nu = 8$ system to a new configuration in which one of the two D-branes has passed through the other and exchanged its relative orientation. Therefore, after this exchange, the total force between the D-branes becomes

$$\begin{aligned}\mathcal{A}' &= \frac{V_1}{2\pi} (8\pi^2\alpha')^{-\frac{1}{2}} \int_0^\infty dt \left(\frac{\pi}{t}\right)^{\frac{1}{2}} e^{-\Delta Y^2/(2\alpha' t)} \left[\left(\frac{f_4}{f_2}\right)^8 - \left(\frac{f_3}{f_2}\right)^8 - 1 \right] \\ &= \frac{V_1}{2\pi\alpha'} |\Delta Y| .\end{aligned}\tag{3.29}$$

This force can be cancelled and the BPS condition restored only if a fundamental string with tension $1/(2\pi\alpha')$ is created when one D-brane passes through another in a $\nu = 8$ system. Notice that, according to this argument, this string creation is an effect of the change in the sign of the R-R contribution due to a parity transformation and *not* to a charge conjugation which would transform one brane into an anti-brane [7]. As a matter of fact, in the case of maximally orthogonal branes, both transformations lead to the same result and are equivalent because the parity preserving part of the R-R sector vanishes. However, this does not happen in general when the two D-branes are tilted with respect to each other [20, 19] since in this case, both the parity preserving and the parity violating parts are non-zero. Finally, as we have explicitly shown, the parity violating part of the R-R force is non-vanishing only when there are $\nu = 8$ directions of mixed type; therefore, in all other cases there is no fundamental string creation when the D-branes pass through each other. Another physical situation, where this parity violating contribution is not trivial, is the magnetic interaction between a D-brane and a dyon, recently examined in [21].

4 Asymmetric BRST invariant R-R states

Like in any string scattering amplitude, also for D-branes one can ask which states mediate the interactions, *i.e.* in our formalism, which states couple to the boundary state and factorize the amplitude $\mathcal{A}$. To answer this question, it is first necessary to discuss in some detail the structure of the closed string states. Since we are ultimately interested in the supergravity interpretation of the D-brane amplitudes, we will limit our considerations to the lowest mass level.

In the NS-NS sector, where no zero-modes are present and no particular technicalities are needed, the structure of the massless states is simple. Nevertheless we find useful to recall here their key properties, since these will be shared also by the R-R states. As shown in Section 2, the boundary state $|B\rangle_{\text{NS}}$ is in the $(-1, -1)$ picture of the NS-NS sector, and thus it can be directly saturated with the usual symmetric states created by the vertex operators

$$V_{\text{NS}}(k; z, \bar{z}) = \epsilon_{\mu\nu} V_{-1}^{\mu}(k/2; z) \tilde{V}_{-1}^{\nu}(k/2; \bar{z}) \quad , \quad (4.1)$$

where

$$V_{-1}^{\mu}(k; z) = c(z) \psi^{\mu}(z) e^{-\phi(z)} e^{ik \cdot X(z)} \quad , \quad (4.2)$$

and the form of $\epsilon_{\mu\nu}$ depends on the particular NS-NS field considered; for example a graviton requires a symmetric and traceless polarization tensor.

The vertex operator (4.1) describes a physical NS-NS state only if it is BRST invariant. To see what are the consequences of this requirement, we split the BRST charge into three parts: $Q = Q_0 + Q_1 + Q_2$ where

$$Q_0 = \oint \frac{dz}{2\pi i} c(z) [T_X(z) + T_{\psi}(z) + T_{\beta\gamma}(z) + \partial c(z)b(z)] \quad , \quad (4.3)$$

$$Q_1 = \frac{1}{2} \oint \frac{dz}{2\pi i} e^{\phi(z)} \eta(z) \psi(z) \cdot \partial X(z) \quad , \quad Q_2 = \frac{1}{4} \oint \frac{dz}{2\pi i} b(z) \eta(z) \partial \eta(z) e^{2\phi(z)} \quad .$$

The first term contains a ghost part and the energy-momentum tensor T of the various fields, the second term is proportional to the supersymmetry current, while the last one contains only ghost and superghost fields. Then, it is not difficult to see that

$$[Q_0 + \tilde{Q}_0, V_{\text{NS}}(k; z, \bar{z})] \propto k^2 (\partial c(z) + \bar{\partial} \tilde{c}(z)) V_{\text{NS}}(k; z, \bar{z}) \quad , \quad (4.4)$$

$$[Q_1 + \tilde{Q}_1, V_{\text{NS}}(k; z, \bar{z})] = 0 \quad \Leftrightarrow \quad k^{\mu} \epsilon_{\mu\nu} = 0 \quad , \quad (4.5)$$

$$[Q_2 + \tilde{Q}_2, V_{\text{NS}}(k; z, \bar{z})] = 0 \quad . \quad (4.6)$$

The vertex (4.1) is therefore BRST invariant only if $k^2 = 0$ and $k \cdot \epsilon = 0$. These two conditions are really *independent* of each other: they originate from two different terms of the BRST charge and can be imposed separately. This means that we can continue off shell the vertex operator (4.1), maintaining Eqs. (4.5)-(4.6) and breaking the BRST invariance only in a soft way, because, whenever $k^2 \neq 0$, the commutator with $Q_0 + \tilde{Q}_0$ remains proportional to the original vertex.

As is well known, perturbative string theory is consistent only if it is BRST invariant. However, a propagating closed string (*i.e.* with $k^{\mu} \neq 0$) emitted from a disk or a D-brane, has to be off shell. In fact, the Neumann boundary conditions along the time direction impose $k_0 = 0$, and thus $k^2 \neq 0$. We believe that, by breaking the BRST invariance in a soft way, *i.e.* by allowing only a non-vanishing commutator with $Q_0 + \tilde{Q}_0$ as in Eq. (4.4), it is still possible to obtain meaningful

results in the field theory limit. Indeed, in Ref. [8], we were able to derive the correct coupling and the correct large distance behavior of the various massless fields of a D-brane solution by simply saturating the boundary state with off-shell closed string states of the lowest mass level.

What we have called soft BRST breaking can be seen as a variant of the off-shell extension proposed in Ref. [16]. The basic idea is that off-shell vertices satisfying Eqs. (4.4)-(4.6) are representatives of cohomology classes of the operator $Q' + \tilde{Q}'$ where

$$Q' = Q - c_0 L_0 - b_0 M \quad , \quad M = -2 \left(\sum_{n>0} n c_{-n} c_n + \sum_{m \geq 0} \gamma_{-m} \gamma_m \right) \quad , \quad (4.7)$$

and similarly for the right moving part. Since

$$(Q' + \tilde{Q}')^2 = -\frac{1}{2} \left[(L_0 + \tilde{L}_0)(M + \tilde{M}) + (L_0 - \tilde{L}_0)(M - \tilde{M}) \right] \quad , \quad (4.8)$$

the operator $Q' + \tilde{Q}'$ can be seen as a BRST charge if it acts either on the usual on-shell states with $L_0 = \tilde{L}_0 = 0$, or on off-shell states such that $L_0 - \tilde{L}_0 = 0$ and $M + \tilde{M} = 0$. The latter conditions are precisely satisfied by the states we are considering. Put differently, the propagating states emitted by D-branes or exchanged in their interactions are off-shell representatives of the cohomology of the restricted charge $Q' + \tilde{Q}'$. Indeed, the ghost zero-modes that must be inserted in Eq. (3.2) break the full BRST invariance, but not the restricted one.

Let us now turn to the R-R sector and, in particular, focus our attention on the massless states of the type IIA theory (of course our results are valid also in the type IIB theory, as discussed in Appendix A). The boundary state $|B\rangle_R$ described in Section 2 is in the $(-1/2, -3/2)$ picture of the R-R sector, and thus, to soak up the superghost number anomaly, it can only couple to states that are also in the asymmetric $(-1/2, -3/2)$ picture. On the contrary, the R-R states that are usually considered in the literature are in the symmetric $(-1/2, -1/2)$ picture and are created by the following vertex operators

$$V_R(k; z, \bar{z}) = \frac{1}{2\sqrt{2}} (CF^{(m+1)})_{\alpha\dot{\beta}} V_{-1/2}^\alpha(k/2; z) \tilde{V}_{-1/2}^{\dot{\beta}}(k/2; \bar{z}) \quad , \quad (4.9)$$

where

$$(CF^{(m+1)})_{\alpha\dot{\beta}} = \frac{(C\Gamma^{\mu_1 \dots \mu_{m+1}})_{\alpha\dot{\beta}}}{(m+1)!} F_{\mu_1 \dots \mu_{m+1}} \quad (4.10)$$

with m odd, and

$$V_\ell^\alpha(k; z) = c(z) S^\alpha(z) e^{\ell \phi(z)} e^{ik \cdot X(z)} \quad . \quad (4.11)$$

The form $F^{(m+1)}$ has the right degree to be interpreted as a R-R field strength of the type IIA theory; indeed the vertex V_R is BRST invariant only if $k^2 = 0$, and $dF^{(m+1)} = d * F^{(m+1)} = 0$ which are precisely the Bianchi and Maxwell equations of a field strength.

To explicitly construct the vertex operators in the $(-1/2, -3/2)$ picture, we first require that they transform into the symmetric vertices (4.9) with a picture changing operation in the right sector [24]. The first guess is [15]

$$W^{(0)}(k; z, \bar{z}) = (CA^{(m)})_{\alpha\beta} V_{-1/2}^\alpha(k/2; z) \tilde{V}_{-3/2}^\beta(k/2; \bar{z}) \quad , \quad (4.12)$$

where $A^{(m)}$ is given by an expression similar to Eq. (4.10). Note that the vertex (4.12) satisfies the same Type IIA GSO projection as V_R , and its polarization contains a form $A^{(m)}$ which has the right degree to be identified with a R-R potential. One can easily check that $W^{(0)}$ commutes with Q_2 , $\tilde{Q}_2$ and $\tilde{Q}_1$, while the commutation with Q_1 yields

$$[Q_1, W^{(0)}(k; z, \bar{z})] = \frac{1}{4\sqrt{2}} (C \not{k} A^{(m)})_{\dot{\alpha}\beta} \eta(z) V_{+1/2}^{\dot{\alpha}}(k/2; z) \tilde{V}_{-3/2}^\beta(k/2; \bar{z}) \quad . \quad (4.13)$$

The right hand side vanishes only if $dA^{(m)} = d * A^{(m)} = 0$. These conditions together imply $k^2 = 0$, and thus guarantee that $W^{(0)}$ commutes also with $Q_0 + \tilde{Q}_0$. Thus, the vertex (4.12) can be made BRST invariant, but then it describes only a pure gauge potential. Moreover, since the two linear conditions coming from Eq. (4.13) imply $k^2 = 0$, the BRST invariance is badly broken when $W^{(0)}$ is naïvely extended off-shell.

In order to describe non trivial gauge potentials in the asymmetric picture of the R-R sector, in Ref. [8] we proposed a generalization of Eq. (4.12). However, the vertex operators constructed there contain explicitly the bare field ξ , and thus do not belong to the (β, γ) system, as it is clear from the bosonization formulas (2.33). Another possibility to describe potentials with non vanishing field strengths, is given by the following vertex operator⁴

$$W_G(k; z, \bar{z}) = \frac{1}{2} [C(A^{(m)} + A^{(m+2)})]_{\alpha\beta} V_{-1/2}^\alpha(k/2; z) \tilde{V}_{-3/2}^\beta(k/2; \bar{z}) \quad . \quad (4.14)$$

Again, W_G is invariant under Q_2 , $\tilde{Q}_2$ and $\tilde{Q}_1$, while the commutation with the linear part of the left BRST charge now yields

$$[Q_1, W_G(k; z, \bar{z})] = \frac{1}{8\sqrt{2}} [C \not{k} (A^{(m)} + A^{(m+2)})]_{\dot{\alpha}\beta} \eta(z) V_{+1/2}^{\dot{\alpha}}(k/2; z) \tilde{V}_{-3/2}^\beta(k/2; \bar{z}) \quad . \quad (4.15)$$

Using the properties of the Γ matrices, one can see that the right hand side is zero if

$$d * A^{(m)} = 0 \quad , \quad dA^{(m)} + * d * A^{(m+2)} = 0 \quad , \quad dA^{(m+2)} = 0 \quad , \quad (4.16)$$

so that Eq. (4.14) describes a non trivial potential $A^{(m)}$ in the Lorentz gauge and a pure gauge field $A^{(m+2)}$ which decouples from all physical amplitudes. Note that the conditions (4.16) together imply that $k^2 = 0$, so that W_G commutes also with

⁴The overall factor of $1/2$ has been introduced for later convenience.

$Q_0 + \tilde{Q}_0$. Finally, by performing a picture changing in the right sector, one can recover the symmetric vertex operator (4.9) with $F^{(m+1)} = dA^{(m)}$. Hence, W_G satisfies all the requirements to be an acceptable R-R vertex operator of the closed string. Just like V_R , also the asymmetric vertex (4.14) has definite left and right superghost numbers, namely $G_0 = \tilde{G}_0 = 0$, even though the only condition that is really necessary in closed string theory is $G_0 + \tilde{G}_0 = 0$. On the other hand, contrarily to what usually happens with the other vertex operators, the polarization of W_G is the sum of two different terms corresponding to two different fields. Moreover, since the constraints for the commutation of W_G with $Q_1 + \tilde{Q}_1$ imply the mass-shell condition $k^2 = 0$, again the BRST invariance is badly broken when the vertex (4.14) is extended off-shell. Thus, W_G commutes with the restricted charge $Q' + \tilde{Q}'$ only if $k^2 = 0$, and hence cannot represent the off-shell states that mediate the interactions between D-branes.

However, there exists yet another possibility to write a non trivial vertex operator in the asymmetric picture. In fact, let us consider again Eq. (4.12) with $dA^{(m)} \neq 0$. To cancel the right hand side of Eq. (4.13), instead of introducing a new gauge potential as in Eq. (4.14), we add to $W^{(0)}$ the following vertex

$$W^{(1)}(k; z, \bar{z}) = -(CA^{(m)})_{\dot{\alpha}\dot{\beta}} \eta(z) V_{+1/2}^{\dot{\alpha}}(k/2; z) \bar{\partial}\tilde{\xi}(\bar{z}) \tilde{V}_{-5/2}^{\dot{\beta}}(k/2; \bar{z}) \quad , \quad (4.17)$$

which commutes with Q_2 and $\tilde{Q}_2$, but not with $\tilde{Q}_1$. Indeed, one finds

$$[\tilde{Q}_1, W^{(1)}(k; z, \bar{z})] = \frac{1}{4\sqrt{2}} (CA^{(m)} \not{k})_{\dot{\alpha}\dot{\beta}} \eta(z) V_{+1/2}^{\dot{\alpha}}(k/2; z) \tilde{V}_{-3/2}^{\dot{\beta}}(k/2; \bar{z}) \quad . \quad (4.18)$$

The right hand sides of Eqs. (4.13) and (4.18) have the same structure and can compensate each other if

$$\not{k}A^{(m)} + A^{(m)}\not{k} = 0 \quad , \quad (4.19)$$

that is, if the potential satisfies the Lorentz gauge condition $d * A^{(m)} = 0$. However, since the commutator of Q_1 with $W^{(1)}$ is not zero, we must add another term $W^{(2)}$ to repair the BRST invariance. The details of this iterative construction are given in Appendix A, and the final result is a vertex operator like

$$W(k; z, \bar{z}) = \sum_{M=0}^{\infty} W^{(M)}(k; z, \bar{z}) \quad , \quad (4.20)$$

where the infinite terms are recursively determined by asking that the commutator of Q_1 with $W^{(M)}$ is canceled by the commutator of $\tilde{Q}_1$ with $W^{(M+1)}$ if Eq. (4.19) is satisfied. This construction ensures that W commutes with $Q_1 + \tilde{Q}_1$ if just the Lorentz gauge is imposed without requiring the mass-shell constraint $k^2 = 0$. The latter is only needed to make W invariant also under Q_0 and $\tilde{Q}_0$. Thus, contrarily to W_G , the new vertex operator W can be extended off-shell by breaking the BRST invariance in a soft way. For this reason, it can be saturated with the boundary state to obtain the correct coupling between a D-brane and a R-R potential. This

property is related to the fact that the off-shell vertices W are representatives of the cohomology classes of the charge $Q' + \tilde{Q}'$ (see Eq. (4.7)), where the superghosts zero-modes have not been singled out contrarily to what suggested in Ref. [16].

Even if the vertex operator (4.20) looks rather complicated, the state that it creates has, instead, a rather simple expression when written in the (β, γ) system. In fact, as shown in detail in Appendix A, we have

$$\begin{aligned} |W\rangle &= \lim_{z, \bar{z} \rightarrow 0} W(k; z, \bar{z}) |0\rangle \\ &= \left(CA^{(m)} \right)_{\alpha\beta} \cos(\gamma_0 \tilde{\beta}_0) |\alpha; k/2\rangle_{-1/2} |\tilde{\beta}; k/2\rangle_{-3/2} \\ &\quad + \left(CA^{(m)} \right)_{\dot{\alpha}\dot{\beta}} \sin(\gamma_0 \tilde{\beta}_0) |\dot{\alpha}; k/2\rangle_{-1/2} |\tilde{\beta}; k/2\rangle_{-3/2} \ , \end{aligned} \quad (4.21)$$

where we have introduced the notation

$$|\alpha; k\rangle_\ell \equiv \lim_{z \rightarrow 0} V_\ell^\alpha(k; z) |0\rangle \ . \quad (4.22)$$

The state $|W\rangle$ is similar in form to the zero-mode part of the boundary state $|B\rangle_R$ (see Eq. (2.30)), and like the latter, is an eigenstate of the total superghost number $G_0 + \tilde{G}_0$ with eigenvalue zero, even though the left and right numbers are not separately well-defined. Asymmetric R-R states with this property have been considered also in Ref. [22]. Despite their very different structure, the two on-shell states $|W\rangle$ of Eq. (4.21) and $|W_G\rangle$, created by the vertex operator (4.14), describe the same physical content and correspond to two different gauge choices for the R-R potentials. More precisely, as we show in Appendix B, $|W\rangle$ and $|W_G\rangle$ are in the same BRST cohomology class, that is⁵

$$|W\rangle = |W_G\rangle + (Q + \tilde{Q}) |\Lambda\rangle \ . \quad (4.23)$$

Note that the BRST equivalence (4.23) guarantees that $|W\rangle$ has a well-defined norm, even if it is created by a vertex operator that contains an infinite number of terms. In fact, as we discuss at length in the next section, there exists a well-defined scalar product for the states $|W\rangle$ which utilizes the same regulator $\mathcal{R}(x)$ appearing in the scalar product of boundary states. Finally, we would like to comment that the asymmetric $(-1/2, -3/2)$ picture of the R-R sector does not necessarily require that the states are proportional to the R-R gauge potentials. For example, the state [22]

$$\left(CF^{(m+1)} \right)_{\alpha\dot{\beta}} (c_0 + \tilde{c}_0) \tilde{\beta}_0 |\alpha; k/2\rangle_{-1/2} |\tilde{\beta}; k/2\rangle_{-3/2} \quad (4.24)$$

is BRST invariant if $F^{(m+1)}$ satisfies the Maxwell equations for a field strength, and transforms to a symmetric state in the $(-1/2, -1/2)$ picture with a picture changing operation in the right sector. In this transformation only the algebraic

⁵For this relation to hold, it is crucial to have the factor of $1/2$ in front of the vertex operator (4.14) that creates $|W_G\rangle$.

part of the BRST charge, (*i.e.* $\tilde{Q}_2$), plays a non trivial role, contrarily to what happens with the states $|W\rangle$, where it is $\tilde{Q}_1$ that acts non trivially and transforms the potential into a field strength. However, states like (4.24) do not play any role in our discussion since, having a non zero total superghost number, they cannot couple to the boundary state.

In the following section we demonstrate that the states $|W\rangle$ correctly factorize the R-R amplitude between two D-branes, and use this result to explain the Coulomb like interaction of the $\nu = 8$ systems from a field theory point of view.

5 Factorization and supergravity analysis of the $\nu = 8$ systems

From the point of view of supergravity the results of Section 3 may appear surprising: in fact, while the absence of force between two parallel D-branes ($\nu = 0$) can be seen as the result of the cancellation among three Feynman diagrams (*i.e.* the exchange of a graviton, a dilaton and a $(p + 1)$ -form potential), it does not seem possible to give a similar interpretation to the no-force condition found for the $\nu = 8$ configurations. In this case the string calculation tells us that the total force due to the exchange of NS-NS states is repulsive and independent of the distance between the two D-branes; on the other hand, even if the two D-branes carry R-R charges of different type, there is a non-vanishing contribution from the R-R sector that exactly cancels the repulsive NS-NS force. In this section we show that there is a very simple interpretation of such a R-R force also from the field theory point of view. For definiteness we will consider in detail the system of one D0 and one D8-brane, but of course our conclusions are valid as well in all other $\nu = 8$ systems related to it by T-duality. The crucial point is the following: for the $\nu = 8$ systems, the charges of the two D-branes are essentially identified, produce the same R-R potential, and therefore, just like in the $\nu = 0$ systems, the R-R interaction is simply due to the usual Coulomb-like force between the D-branes. It should not be surprising that forms of different order can be identified. One well-known example of this phenomenon is the electro-magnetic duality of the Type II theories. In fact, let us consider the massless R-R state created by the symmetric vertex operator (4.9)

$$|V\rangle = \frac{(C\Gamma^{\mu_1 \dots \mu_{m+1}})_{\alpha\dot{\beta}}}{2\sqrt{2}(m+1)!} F_{\mu_1 \dots \mu_{m+1}} |\alpha; k/2\rangle_{-1/2} |\tilde{\beta}; k/2\rangle_{-1/2} . \quad (5.1)$$

Due to the structure of the polarization factor in Eq. (5.1), it is easy to see that the field strengths satisfy the duality condition $F^{(m+1)} \simeq *F^{(9-m)}$. Such a relation imposes a duality constraint also on the physical degrees of freedom of the potentials $A^{(m)}$. For instance, if the space momentum of the state (5.1) lies entirely along the 9th direction, the duality condition for a 2-form reads

$$F_{01} = k_0 A_1 = -F_{2\dots 9} = k_9 A_{2\dots 8} . \quad (5.2)$$

Since the state (5.1) is massless, one has also $k_0 = k_9$, and thus

$$A_1 = A_{2\dots 8} \quad . \quad (5.3)$$

In this way one recovers the usual duality relation for the potential fields $A^{(m)} \simeq *A^{(8-m)}$ in the transverse space. It is important to realize that this relation involves only the physical transverse degrees of freedom of the potentials, and does not hold for the non-physical polarizations that are the analogues of the scalar and longitudinal photons of electrodynamics. Usually, such states are never taken into account in string theory because they decouple from any physical amplitude. However, as we shall see, they are particularly relevant to the present discussion because a mixture of longitudinal and scalar states provides a local description of the Coulomb force between D-branes, which may appear as an instantaneous effective interaction when just the physical degrees of freedom are quantized. Note that, if only closed strings are present, there are no R-R charged objects, and thus the longitudinal and scalar polarizations do not appear as propagating states either. Even if these degrees of freedom always decouple, we can still write some states that describe them in a pure closed string framework. These are created by the vertex operator (4.12), and are explicitly given by

$$|W^{(0)}\rangle = \frac{1}{m!} (C\Gamma^{\mu_1\dots\mu_m})_{\alpha\beta} A_{\mu_1\dots\mu_m} |\alpha; k/2\rangle_{-1/2} |\tilde{\beta}; k/2\rangle_{-3/2} \quad . \quad (5.4)$$

In fact, as we discussed in Section 4, $|W^{(0)}\rangle$ is BRST invariant if $k^2 = 0$ and $dA^{(m)} = d * A^{(m)} = 0$ which are precisely the conditions fulfilled by the mixture of longitudinal and scalar polarizations that describes the Coulomb interaction. For example a 1-form potential of this type with $k_0 = k_9$ is characterized by

$$A_0 = A_9 \quad , \quad \text{and} \quad A_i = 0 \quad \text{for} \quad i = 1, \dots, 8 \quad . \quad (5.5)$$

The spinorial structure of Eq. (5.4) implies that the unphysical polarizations satisfy a 10-dimensional Hodge duality similar to the one of the field strength; for instance, in the case of a 1-form we have $A_9 = -A_{01\dots 8}$, which combined with Eq. (5.5) leads to the following relation

$$A_0 = -A_{01\dots 8} \quad . \quad (5.6)$$

The two dualities (5.3) and (5.6), and their obvious generalizations, can be unified by saying that the components of two potentials are identified if they have the same longitudinal indices and their *transverse* indices are complementary. The unusual relation (5.6) is of no relevance in perturbative string theory where the unphysical degrees of freedom always decouple, but it becomes important when dealing with boundary states. In this case it has remarkable consequences: in fact, it implies that the charge felt by A_0 is opposite to the charge felt by $A_{01\dots 8}$, and thus the attractive Coulomb R-R force between a D0 and a D8 brane can be interpreted as due to the exchange of longitudinal and scalar polarizations identified according

to Eq. (5.6). Note that this argument is consistent with the results of Section 3, where we have found that, in $\nu = 8$ systems, the R-R force does not receive any correction from the massive string states and is exactly the opposite of the force due to the exchange of massless R-R fields between two parallel D-branes. To see that this interpretation is correct, we actually need to verify that the states propagating between two D-branes satisfy duality relations like (5.6). Since these states are off-shell, they can not be like $|W^{(0)}\rangle$ of Eq. (5.4), but instead are like $|W\rangle$ of Eq. (4.21). Thus, henceforth we focus on asymmetric states of this type and study the Hilbert space they generate. As a first step, we define the conjugated state $\langle W|$ which satisfies the same GSO projection of $|W\rangle$, namely

$$\begin{aligned} \langle W| = & {}_{-3/2}\langle\dot{\alpha}; k/2| {}_{-1/2}\langle\tilde{\beta}; k/2| \cos(\beta_0\tilde{\gamma}_0) (CA^{(m)})_{\dot{\alpha}\tilde{\beta}} \\ & + {}_{-3/2}\langle\alpha; k/2| {}_{-1/2}\langle\tilde{\beta}; k/2| \sin(\beta_0\tilde{\gamma}_0) (CA^{(m)})_{\alpha\tilde{\beta}} \quad . \end{aligned} \quad (5.7)$$

However, with this definition the naïve scalar product between a bra and a ket is divergent or ill defined due to the infinite contributions of the superghosts, just like the naïve scalar product between two boundary states. We overcome this problem by regularizing the scalar product with the same prescription used in Section 3 for the amplitude between two boundary states. Thus, we define

$$\langle W', W \rangle \equiv \lim_{x \rightarrow 1} \langle W' | \mathcal{R}(x) | W \rangle \quad , \quad (5.8)$$

where the regulator $\mathcal{R}(x)$ is given in Eq. (3.9). It is now easy to see that, with this prescription, the states of Eq. (4.21) have a definite norm. For example, following the same procedure outlined in Section 3, for the case of 1-forms we have

$$\begin{aligned} \langle W, W \rangle &= \lim_{x \rightarrow 1} \left[\frac{1}{2(1+x^2)} \langle A | \langle \tilde{B} | (CA^{(1)})_{AB} x^{2F_0} (CA^{(1)})_{CD} | C \rangle | \tilde{D} \rangle \right. \\ &\quad \left. + \frac{1}{2(1-x^2)} \langle A | \langle \tilde{B} | (CA^{(1)})_{AB} x^{2F_0} (CA^{(1)})_{\Gamma 11} | C \rangle | \tilde{D} \rangle \right] \\ &= 8 A_\mu A^\mu \quad . \end{aligned} \quad (5.9)$$

Note that the second line does not contribute, since the result of the scalar product over the fermionic zero-modes goes to zero faster than $(1-x^2)$ when $x \rightarrow 1$. The factor of 8 in the final result correctly counts the physical degeneracy of the R-R vacuum, and is the product of the superghost contribution (*i.e.* $1/2$) and of the chiral trace over the Γ matrices (*i.e.* 16). Thus, our regularization prescription makes manifest the role of the superghost zero-modes in the R-R sector, that is to halve the degeneracy of the fermionic vacuum. Moreover, with the definition (5.8), the one-to-one correspondence between $|W\rangle$ and $|W_G\rangle$ displayed in Eq. (4.23) becomes an isometry: in fact, the scalar product (5.8) reduces to the usual one when $|W\rangle$ is written in terms of $|W_G\rangle$. But the most striking feature of the scalar product (5.8) is that forms of different order are, in general, not orthogonal to each

other. To see this explicitly, let us consider for example the subspace generated by

$$\begin{aligned} |W_1\rangle &= (C\Gamma^1)_{\alpha\beta} \cos(\gamma_0\tilde{\beta}_0) |\alpha; k/2\rangle_{-1/2} |\tilde{\beta}; k/2\rangle_{-3/2} \\ &+ (C\Gamma^1)_{\dot{\alpha}\dot{\beta}} \sin(\gamma_0\tilde{\beta}_0) |\dot{\alpha}; k/2\rangle_{-1/2} |\dot{\tilde{\beta}}; k/2\rangle_{-3/2} \quad , \end{aligned} \quad (5.10)$$

and

$$\begin{aligned} |W_{2\dots 8}\rangle &= (C\Gamma^{2\dots 8})_{\alpha\beta} \cos(\gamma_0\tilde{\beta}_0) |\alpha; k/2\rangle_{-1/2} |\tilde{\beta}; k/2\rangle_{-3/2} \\ &+ (C\Gamma^{2\dots 8})_{\dot{\alpha}\dot{\beta}} \sin(\gamma_0\tilde{\beta}_0) |\dot{\alpha}; k/2\rangle_{-1/2} |\dot{\tilde{\beta}}; k/2\rangle_{-3/2} \quad , \end{aligned} \quad (5.11)$$

where the momentum lies only in the 0th and 9th directions in order to keep the transversality condition. These two states are not perpendicular to each other, because in the mixed scalar product the analogue of the second line of Eq. (5.9) gives a non-vanishing contribution leading to $\langle W_1, W_{2\dots 8} \rangle = 8$. We then conclude that, in the subspace spanned by $|W_1\rangle$ and $|W_{2\dots 8}\rangle$, the scalar product (5.8) defines a degenerate metric proportional to

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \quad (5.12)$$

with a null-state

$$|\zeta\rangle = |W_1\rangle - |W_{2\dots 8}\rangle \quad , \quad (5.13)$$

which decouples from all amplitudes, even if boundary states are present. Thus, we can set $|\zeta\rangle = 0$, and identify the components of a R-R field along $|W_1\rangle$ and $|W_{2\dots 8}\rangle$, recovering in this way the off-shell extension of the duality relation (5.3). These arguments hold for all states with transverse polarizations, while for longitudinal and scalar states the results are slightly different, even if the analysis is similar. Let us consider, for example, the 1-form state that carries the Coulomb interaction (5.5), and continues it off-shell in the kinematic region that is relevant for the study of D-brane interactions, namely

$$\begin{aligned} |W_0\rangle &= (C\Gamma^0)_{\alpha\beta} \cos(\gamma_0\tilde{\beta}_0) |\alpha; k/2\rangle_{-1/2} |\tilde{\beta}; k/2\rangle_{-3/2} \\ &+ (C\Gamma^0)_{\dot{\alpha}\dot{\beta}} \sin(\gamma_0\tilde{\beta}_0) |\dot{\alpha}; k/2\rangle_{-1/2} |\dot{\tilde{\beta}}; k/2\rangle_{-3/2} \quad , \end{aligned} \quad (5.14)$$

where $k_i = 0$ for $i = 0, \dots, 8$. It is very easy to verify that the following 9-form state

$$\begin{aligned} |W_{0\dots 8}\rangle &= (C\Gamma^{0\dots 8})_{\alpha\beta} \cos(\gamma_0\tilde{\beta}_0) |\alpha; k/2\rangle_{-1/2} |\tilde{\beta}; k/2\rangle_{-3/2} \\ &+ (C\Gamma^{0\dots 8})_{\dot{\alpha}\dot{\beta}} \sin(\gamma_0\tilde{\beta}_0) |\dot{\alpha}; k/2\rangle_{-1/2} |\dot{\tilde{\beta}}; k/2\rangle_{-3/2} \end{aligned} \quad (5.15)$$

is not orthogonal to $|W_0\rangle$. In fact, in the scalar product $\langle W_0, W_{0\dots 8} \rangle$, the two Γ_0 matrices present in both states cancel, and so the calculation becomes identical to

the one outlined for $\langle W_1, W_{2\dots 8} \rangle$. Again, in the subspace spanned by the states (5.14) and (5.15), the scalar product leads to a degenerate metric proportional to

$$\begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \quad (5.16)$$

which admits the null state

$$|\chi\rangle = |W_0\rangle + |W_{0\dots 8}\rangle \quad . \quad (5.17)$$

While on-shell the scalar and longitudinal polarizations like (5.5) always decouple from physical amplitudes, the presence of a D-brane forces an off-shell continuation and only linear combinations like (5.17) decouple. When we set $|\chi\rangle = 0$, we are led to identify A_0 with $-A_{0\dots 8}$, recovering in this way the duality relation (5.6) also off-shell. Thus, the Hilbert space structure is responsible both for the identification (5.3), that is usually seen as an effect of the GSO projection of the symmetrical states, and for the identification (5.6), that is proper only of the longitudinal asymmetric states (4.21), since these have no symmetric counterparts.

At the beginning of this section, we suggested that the linear R-R potential in the $\nu = 8$ systems is simply the analogue of the usual Coulomb electric potential between two charges of opposite sign; this is because the two D-branes produce essentially the same R-R field, even if their charges seem different. Now we can explicitly prove that this interpretation is correct by identifying the R-R state exchanged in the interaction between the two D-branes. In order to exploit the explicit formulas that we have derived before, we consider a system of one 0-brane and one 8-brane, but it is clear that this analysis is general and can be applied to all $\nu = 8$ configurations. Focusing on the massless sector, the R-R amplitude between the two boundary states, $\mathcal{A}_{R-R}^0$, can be factorized by writing the identity operator with the asymmetric states of Eq. (4.21). In particular the relevant terms are

$$\mathbb{1} = |W_0\rangle \langle W_0| + P(|\chi\rangle) + \dots \quad , \quad (5.18)$$

where P is a projector onto the one dimensional space generated by the vector $|\chi\rangle$. From our previous analysis of the R-R Hilbert space structure, we know that the combination $|\chi\rangle$ can be ignored because it always decouples, while the 1-form state $|W_0\rangle$ has a non-zero overlapping with both the 0-brane and the 8-brane. Inserting twice the identity (5.18) in the Eq. (3.2), it is easy to see that the massless R-R contribution to the D-brane interaction can be written as

$$\mathcal{A}_{R-R}^0 = {}_R\langle B_0 | W_0 \rangle \langle W_0 | D | W_0 \rangle \langle W_0 | B_8 \rangle_R \quad , \quad (5.19)$$

where, henceforth, the state $|W_0\rangle$ is normalized to one⁶, while all the other R-R states that give no contribution are understood. With this normalization one can

⁶This simply amounts to multiply Eq. (5.14) by $1/(2\sqrt{2})$.

verify that, in the field theory limit, the scattering amplitude among the asymmetric R-R states and the NS-NS ones are correctly reproduced by the following action

$$S = \int d^{10}x \sqrt{-g} \left[-\frac{1}{2(m+1)!} e^{\frac{5-(m+1)}{\sqrt{2}}\kappa\phi} \left(F_{\mu_1 \dots \mu_{m+1}}^{(m+1)} \right)^2 \right] , \quad (5.20)$$

where, as usual, $F^{(m+1)} = dA^{(m)}$.

We can now reinterpret the result of the factorization (5.19) as the missing Feynman diagram describing the R-R interaction from a field theory point of view. In fact, the closed string propagator D in Eq. (5.19) simply becomes the usual field theory propagator in the Feynman gauge $-1/k^2$, that is consistent with the action (5.20). On the other hand, from Ref. [8], we know that, when a boundary state is directly saturated with a closed string state, one gets the field theory vertices that are usually derived from the Born-Infeld lagrangian, namely

$$\langle W_0 | B_0 \rangle_R = -\langle W_0 | B_8 \rangle_R = -\sqrt{2}T_0 V_1 . \quad (5.21)$$

The coupling of a D8-brane and a 1-form potential is not manifest at the level of the D-brane action, because it is an effect of the duality relation (5.6). As is well known for the type IIB theory and other similar cases, it is rather difficult to implement a duality relation directly in an action, and normally infinite auxiliary fields are needed for this purpose.

Appendix A

In this appendix we prove the BRST invariance of the vertex operator

$$W(k; z, \bar{z}) = \sum_{M=0}^{\infty} W^{(M)}(k; z, \bar{z}) \quad (A.1)$$

introduced in Section 4 to describe the emission of a R-R field in the asymmetric $(-1/2, -3/2)$ picture. We treat simultaneously Type IIA and Type IIB theories assuming that the GSO projection is defined in Eq. (2.27) (with p even for Type IIA, and p odd for Type IIB), and use 32-dimensional spinor indices. For notational convenience, we introduce also the chiral projectors $\Pi_q \equiv (1 + (-1)^q \Gamma_{11})/2$, and the following combinations:

$$\begin{aligned} \mathcal{V}_{-1/2+M}^A(z) &= \partial^{M-1} \eta(z) \dots \eta(z) c(z) S^A(z) e^{(-\frac{1}{2}+M)\phi(z)} e^{ik \cdot X(z)/2} , \\ \tilde{\mathcal{V}}_{-3/2-M}^A(\bar{z}) &= \bar{\partial}^M \tilde{\xi}(\bar{z}) \dots \bar{\partial} \tilde{\xi}(\bar{z}) \tilde{c}(\bar{z}) \tilde{S}^A(\bar{z}) e^{(-\frac{3}{2}-M)\tilde{\phi}(\bar{z})} e^{ik \cdot \tilde{X}(\bar{z})/2} , \end{aligned} \quad (A.2)$$

where S^A are the spin fields and ϕ, ξ and η come from the superghost fermionization (see Eq. (2.33)). With these notations and dropping for simplicity the dependence

on the momentum k , all terms in Eq. (A.1) can be written in a compact way as

$$W^{(M)}(z, \bar{z}) = a_M \left[\Pi_{p+M} C A^{(m)} \Pi_M \right]_{AB} \mathcal{V}_{-1/2+M}^A(z) \tilde{\mathcal{V}}_{-3/2-M}^B(\bar{z}) \quad , \quad (\text{A.3})$$

where $a_0 = 1$ and the other coefficients a_M will be specified later. Converting to the Majorana-Weyl notation, one can easily recognize for instance that Eq. (A.3) for $M = 0$ in a Type IIA theory (p even and m odd) reads

$$\left(C A^{(m)} \right)_{\alpha\beta} c(z) S^\alpha(z) e^{-\phi(z)/2} e^{ik \cdot X(z)/2} \tilde{c}(\bar{z}) \tilde{S}^\beta(\bar{z}) e^{-3\tilde{\phi}(\bar{z})/2} e^{ik \cdot \tilde{X}(\bar{z})/2} \quad , \quad (\text{A.4})$$

which is precisely Eq. (4.12). Similarly, one can check that Eq. (A.3) for $M = 1$ reduces to Eq. (4.17) if $a_1 = 1$. We now study the BRST properties of the vertex W , and to do this we split the BRST charge according to Eq. (4.3). It is well known that any operator with conformal dimension 0 and of the type $c(z)U(z)$, where $U(z)$ is a primary field made up only by matter and superghost fields, commutes with Q_0 . Then, it follows that the commutation with $Q_0 + \tilde{Q}_0$ is equivalent to the on-shell condition

$$\left[Q_0, W^{(M)}(z, \bar{z}) \right] = \left[\tilde{Q}_0, W^{(M)}(z, \bar{z}) \right] = 0 \Leftrightarrow k^2 = 0 \quad . \quad (\text{A.5})$$

It is also easy to show, by direct computation of the relevant OPEs, that $W^{(M)}$ commutes with Q_2 and $\tilde{Q}_2$. The only non-trivial commutators are the ones with Q_1 and $\tilde{Q}_1$. To evaluate them, we make use of the following OPEs [23]

$$(\psi^\mu)_1(w) S_{-n/2}^A(z) \sim \frac{i}{\sqrt{2}} (w-z)^{\frac{n-1}{2}} (\Gamma_{11} \Gamma^\mu)_B^A S_{-n/2+1}^B(z) \quad , \quad (\text{A.6})$$

where we abbreviated $(\psi_\mu)_1(z) \equiv \psi_\mu(z) e^{\phi(z)}$ and $S_{-n/2}^A(z) \equiv S^A(z) e^{-\frac{n}{2}\phi(z)}$, and

$$\partial X^\mu(w) e^{ik \cdot X(z)/2} \sim \frac{i(k^\mu/2) e^{ik \cdot X(z)/2}}{w-z} \quad , \quad (\text{A.7})$$

plus of course their tilded counterparts. Straightforwardly enough one obtains

$$\begin{aligned} \left[Q_1, W^{(M)}(z, \bar{z}) \right] &= \frac{a_M (-1)^{p+M}}{4\sqrt{2}M!} \left(\Pi_{p+M+1} C \not{k} A^{(m)} \Pi_M \right)_{AB} \\ &\times \mathcal{V}_{-\frac{1}{2}+(M+1)}^A(z) \tilde{\mathcal{V}}_{-\frac{3}{2}-M}^B(\bar{z}) \quad , \end{aligned} \quad (\text{A.8})$$

$$\begin{aligned} \left[\tilde{Q}_1, W^{(M+1)}(z, \bar{z}) \right] &= \frac{-a_{M+1} (M+1)!}{4\sqrt{2}} \left(\Pi_{p+M+1} C A^{(m)} \not{k} \Pi_M \right)_{AB} \\ &\times \mathcal{V}_{-\frac{1}{2}+(M+1)}^A(z) \tilde{\mathcal{V}}_{-\frac{3}{2}-M}^B(\bar{z}) \quad . \end{aligned} \quad (\text{A.9})$$

Therefore, we have

$$\left[Q_1, W^{(M)}(z, \bar{z}) \right] + \left[\tilde{Q}_1, W^{(M+1)}(z, \bar{z}) \right] = 0 \quad , \quad (\text{A.10})$$

provided the coefficients a_M are given by

$$a_M = \frac{(-)^{\frac{1}{2}M(M+1)}}{M![(M-1)!\dots 2\cdot 1]^2} , \quad (\text{A.11})$$

and the R-R gauge field satisfies

$$(-)^p \not{A} A^{(m)} + A^{(m)} \not{A} = 0 , \quad (\text{A.12})$$

that is if the potential is in the generalized Lorentz gauge $d * A^{(m)} = 0$. If this is the case, and recalling that

$$[\tilde{Q}_1, W^{(0)}(z, \bar{z})] = 0 , \quad (\text{A.13})$$

it follows that the massless vertex (A.1) is BRST invariant,

$$[Q + \tilde{Q}, W(z, \bar{z})] = 0 . \quad (\text{A.14})$$

We now show that under a picture changing operation in the right sector [24],

$$W(z, \bar{z}) \rightarrow [Q + \tilde{Q}, 2\tilde{\xi}(\bar{z})W(z, \bar{z})]_+ , \quad (\text{A.15})$$

the asymmetric vertex W transforms into the usual operator V_R in the symmetric $(-1/2, -1/2)$ picture whose polarization contains the R-R field strength as shown in Eq. (4.9). To do this, we firstly notice that since the insertion of $\tilde{\xi}$ does not alter the conformal dimension of W , the commutation with $\tilde{Q}_0$ is still ensured if $k^2 = 0$; secondly, also the commutation with $\tilde{Q}_2$ is not spoiled. Again, the non-trivial part is the commutator with $\tilde{Q}_1$. Utilizing the OPEs (A.6, A.7), we can see that only the first term $W^{(0)}$ is responsible for the picture-changing; indeed, we get

$$\begin{aligned} [\tilde{Q}_1, 2\tilde{\xi}(\bar{z})W^{(0)}(z, \bar{z})]_+ &= \frac{1}{2\sqrt{2}} (\Pi_p C A^{(m)} \not{A} \Pi_1)_{AB} \mathcal{V}_{-\frac{1}{2}}^A(z) \tilde{\mathcal{V}}_{-\frac{1}{2}}^B(\bar{z}) \\ &= V_R(z, \bar{z}) , \end{aligned} \quad (\text{A.16})$$

where in the last step we used the Lorentz gauge condition. For the remaining terms $W^{(M)}$ with $M > 0$, the only non-zero contribution to the picture-changing arises when $\tilde{\xi}$ acts as a spectator, so that

$$[\tilde{Q}_1, 2\tilde{\xi}(\bar{z})W^{(M)}(z, \bar{z})]_+ = -2\tilde{\xi}(\bar{z}) [\tilde{Q}_1, W^{(M)}(z, \bar{z})] . \quad (\text{A.17})$$

Thus, taking into account Eqs. (A.13) and (A.14), we find that

$$[Q + \tilde{Q}, 2\tilde{\xi}(\bar{z})W(z, \bar{z})]_+ = V_R(z, \bar{z}) - 2\tilde{\xi}(\bar{z}) [Q + \tilde{Q}, W(z, \bar{z})] = V_R(z, \bar{z}) . \quad (\text{A.18})$$

We conclude this appendix by finding the explicit expression of the state $|W\rangle$ created by the vertex operator (A.1). For the first term, $W^{(0)}$, we have by definition

$$|W^{(0)}\rangle = (\Pi_p C A^{(m)} \Pi_0)_{AB} |A; k/2\rangle_{-1/2} |\tilde{B}; k/2\rangle_{-3/2} . \quad (\text{A.19})$$

Consider now the second term, $W^{(1)}$. We have

$$\begin{aligned} |W^{(1)}\rangle &= - \left(\Pi_{p+1} C \mathcal{A}^{(m)} \Pi_1 \right)_{AB} |A; k/2\rangle |\tilde{B}; k/2\rangle \\ &\times \lim_{z \rightarrow 0} \eta(z) |1/2\rangle_\phi \lim_{\bar{z} \rightarrow 0} \bar{\partial} \tilde{\xi}(\bar{z}) | -5/2\rangle_{\tilde{\phi}} , \end{aligned} \quad (\text{A.20})$$

where the notation $|\ell\rangle_\phi$ stands for the vacuum of the (ϕ, η, ξ) system with ϕ -momentum ℓ , which, according to Eq. (2.35), coincides with the superghost vacuum in the $P = \ell$ picture. Keeping this in mind, and using the fermionization formulas (2.33), it is easy to see that

$$\begin{aligned} \lim_{z \rightarrow 0} \eta(z) |1/2\rangle_\phi &= - \lim_{z \rightarrow 0} z^{-\frac{1}{2}} \gamma(z) |P = -1/2\rangle = - \gamma_0 |P = -1/2\rangle , \\ \lim_{\bar{z} \rightarrow 0} \bar{\partial} \tilde{\xi}(\bar{z}) | -5/2\rangle_{\tilde{\phi}} &= \lim_{\bar{z} \rightarrow 0} \bar{z}^{\frac{3}{2}} \tilde{\beta}(\bar{z}) |\tilde{P} = -3/2\rangle = \tilde{\beta}_0 |\tilde{P} = -3/2\rangle , \end{aligned} \quad (\text{A.21})$$

so that

$$|W^{(1)}\rangle = (\Pi_{p+1} C \mathcal{A}^m \Pi_1)_{AB} \gamma_0 \tilde{\beta}_0 |A; k/2\rangle_{-1/2} |\tilde{B}; k/2\rangle_{-3/2} . \quad (\text{A.22})$$

It is clear that all terms of W with M even (odd) will give exactly the same structure of $|W^{(0)}\rangle$ ($|W^{(1)}\rangle$), except for the superghost part. To deal with it, we use the generalization of the formulas in Eq. (A.21), namely

$$\begin{aligned} \lim_{z \rightarrow 0} \partial^{M-1} \eta(z) \dots \eta(z) | -1/2 + M\rangle_\phi &= (-)^{\frac{M(M+1)}{2}} (M-1)! \dots 2! \gamma_0^M |P = -1/2\rangle , \\ \lim_{\bar{z} \rightarrow 0} \bar{\partial}^M \tilde{\xi}(\bar{z}) \dots \bar{\partial} \tilde{\xi}(\bar{z}) | -3/2 - M\rangle_{\tilde{\phi}} &= (-)^{\frac{M(M-1)}{2}} (M-1)! \dots 2! \tilde{\beta}_0^M |\tilde{P} = -3/2\rangle . \end{aligned} \quad (\text{A.23})$$

Then, using the explicit expression of $W^{(M)}$, Eq. (A.3) and Eq. (A.11), we finally obtain

$$\begin{aligned} |W\rangle &= \left\{ \left(\Pi_p C \mathcal{A}^{(m)} \Pi_0 \right)_{AB} \cos(\gamma_0 \tilde{\beta}_0) + \left(\Pi_{p+1} C \mathcal{A}^{(m)} \Pi_1 \right)_{AB} \sin(\gamma_0 \tilde{\beta}_0) \right\} \\ &\times |A; k/2\rangle_{-1/2} |\tilde{B}; k/2\rangle_{-3/2} . \end{aligned} \quad (\text{A.24})$$

Converting to the Majorana-Weyl notation, one can check that the state (A.24) for a Type IIA theory coincides with Eq. (4.21).

Appendix B

In this appendix we prove the BRST equivalence between the on-shell R-R vertex operators W_G and W defined respectively in Eqs. (4.14) and (4.20). For definiteness, we study in detail vertices associated to a 1-form potential, but of course our results are completely general. Let us then consider the vertex W_G for a 1-form, which explicitly reads

$$W_G(k; z, \bar{z}) = \frac{1}{2} \left[A_\mu (C \Gamma^\mu)_{\alpha\beta} + \frac{1}{3!} A_{\mu\nu\rho} (C \Gamma^{\mu\nu\rho})_{\alpha\beta} \right] \mathcal{V}_{-1/2}^\alpha(z) \tilde{\mathcal{V}}_{-3/2}^\beta(\bar{z}) , \quad (\text{B.1})$$

where $\mathcal{V}_\ell^A$ is defined in Eq. (A.2). This vertex is BRST invariant if (see Eq. (4.16))

$$k^\mu A_\mu = 0 \quad , \quad k_\mu A_\nu - k_\nu A_\mu + k^\rho A_{\rho\mu\nu} = 0 \quad , \quad k_{[\lambda} A_{\mu\nu\rho]} = 0 \quad . \quad (\text{B.2})$$

These conditions imply that $k^2 = 0$, and that the 3-form potential is pure gauge.

We now show that by adding to Eq. (B.1) a BRST exact operator we can reconstruct the vertex W with indefinite left and right superghost numbers. To this aim, let us begin by considering the following operator

$$\Lambda^{(0)}(k; z, \bar{z}) = \sqrt{2} \Lambda_{\mu\nu} (C\Gamma^{\mu\nu})_{\alpha\dot{\beta}} \mathcal{V}_{-1/2}^\alpha(z) \tilde{\mathcal{V}}_{-5/2}^{\dot{\beta}}(\bar{z}) \quad , \quad (\text{B.3})$$

where $k^2 = 0$ and $\Lambda_{\mu\nu}$ is a 2-form to be specified later. It is simple to show that $\Lambda^{(0)}$ commutes with Q_0 , $\tilde{Q}_0$, Q_2 and $\tilde{Q}_2$, whereas the commutators with Q_1 and $\tilde{Q}_1$ are not vanishing and read

$$[Q_1, \Lambda^{(0)}(k; z, \bar{z})] = \frac{1}{4} \Lambda_{\mu\nu} (C \not{k} \Gamma^{\mu\nu})_{\dot{\alpha}\dot{\beta}} \mathcal{V}_{+1/2}^{\dot{\alpha}}(z) \tilde{\mathcal{V}}_{-5/2}^{\dot{\beta}}(\bar{z}) \quad , \quad (\text{B.4})$$

$$[\tilde{Q}_1, \Lambda^{(0)}(k; z, \bar{z})] = \frac{1}{4} \Lambda_{\mu\nu} (C\Gamma^{\mu\nu} \not{k})_{\alpha\beta} \mathcal{V}_{-1/2}^\alpha(z) \tilde{\mathcal{V}}_{-3/2}^\beta(\bar{z}) \quad . \quad (\text{B.5})$$

The right hand side of Eq. (B.5) has the same operator structure of W_G and also of the first term of W , *i.e.* $W^{(0)}$, whereas the right hand side of Eq. (B.4) has the same operator structure of $W^{(1)}$. Thus, we can reach our goal by simply adjusting the polarization coefficients. In fact, if we choose $\Lambda_{\mu\nu}$ such that

$$\Lambda_{\mu\nu} k^\nu = A_\nu \quad , \quad (\text{B.6})$$

it follows that

$$[\tilde{Q}_1, \Lambda^{(0)}(k; z, \bar{z})] = \frac{1}{2} \left[A_\mu (C\Gamma^\mu)_{\alpha\beta} - \frac{1}{3!} A'_{\mu\nu\rho} (C\Gamma^{\mu\nu\rho})_{\alpha\beta} \right] \mathcal{V}_{-1/2}^\alpha(z) \tilde{\mathcal{V}}_{-3/2}^\beta(\bar{z}) \quad , \quad (\text{B.7})$$

where $A'_{\mu\nu\rho} = -k_{[\mu} \Lambda_{\nu\rho]}$. Since this 3-form satisfies exactly the same properties as $A_{\mu\nu\rho}$ in Eq. (B.2), we can identify them so that we have

$$\begin{aligned} W_G(k; z, \bar{z}) + [\tilde{Q}_1, \Lambda^{(0)}(k; z, \bar{z})] &= A_\mu (C\Gamma^\mu)_{\alpha\beta} \mathcal{V}_{-1/2}^\alpha(z) \tilde{\mathcal{V}}_{-3/2}^\beta(\bar{z}) \\ &= W^{(0)}(k; z, \bar{z}) \quad . \end{aligned} \quad (\text{B.8})$$

Let us now consider the following operator

$$\Lambda^{(1)}(k; z, \bar{z}) = -\frac{\sqrt{2}}{2} \Lambda_{\mu\nu} (C\Gamma^{\mu\nu})_{\dot{\alpha}\dot{\beta}} \mathcal{V}_{1/2}^{\dot{\alpha}}(z) \tilde{\mathcal{V}}_{-7/2}^{\dot{\beta}}(\bar{z}) \quad , \quad (\text{B.9})$$

with $k^2 = 0$, which commutes with Q_0 , $\tilde{Q}_0$, Q_2 and $\tilde{Q}_2$. Then, using Eqs. (B.6) and (B.4), straightforwardly enough we get

$$\begin{aligned} [Q_1, \Lambda^{(0)}(k; z, \bar{z})] + [\tilde{Q}_1, \Lambda^{(1)}(k; z, \bar{z})] &= -A_\mu (C\Gamma^\mu)_{\dot{\alpha}\dot{\beta}} \mathcal{V}_{+1/2}^{\dot{\alpha}}(z) \tilde{\mathcal{V}}_{-5/2}^{\dot{\beta}}(\bar{z}) \\ &= W^{(1)}(k; z, \bar{z}) \quad . \end{aligned} \quad (\text{B.10})$$

This procedure can be iterated because it is always possible to find operators $\Lambda^{(M)}(k; z, \bar{z})$ such that

$$\left[Q_1, \Lambda^{(M-1)}(k; z, \bar{z})\right] + \left[\tilde{Q}_1, \Lambda^{(M)}(k; z, \bar{z})\right] = W^{(M)}(k; z, \bar{z}) \quad (\text{B.11})$$

for any $M \geq 0$. Combining all these equations, we finally obtain

$$W_G(k; z, \bar{z}) + \left[Q_1 + \tilde{Q}_1, \Lambda(k; z, \bar{z})\right] = W(k; z, \bar{z}) , \quad (\text{B.12})$$

where $\Lambda(k; z, \bar{z}) = \sum_M \Lambda^{(M)}(k; z, \bar{z})$. This equation implies that the states created by W_G and W are related to each other as shown in Eq. (4.23). The states $|W\rangle$ and $|W_G\rangle$ are BRST equivalent only on shell, while if $k^2 \neq 0$ they are not related even in the restricted cohomology.

References

- [1] J. Polchinski, Phys. Rev. Lett **75** (1995) 4724, [hep-th/9510017](#).
- [2] J. Polchinski, S. Chaudhuri and C.V. Johnson, *Notes on D-branes*, [hep-th/9602052](#); J. Polchinski, *TASI lectures on D-branes*, [hep-th/9611050](#).
- [3] M.B. Green and P. Wai, Nucl. Phys. **B431** (1994) 131; M. Li, Nucl. Phys. **B460** (1996) 351, [hep-th/9510161](#); C.G. Callan and I.R. Klebanov, Nucl. Phys. **B465** (1996) 473, [hep-th/9511173](#); C. Schmidhuber, Nucl. Phys. **B467** (1996) 146, [hep-th/9601003](#); M.B. Green and M. Gutperle, Nucl. Phys. **B476** (1996) 484, [hep-th/9604091](#).
- [4] M. Billó, D. Cangemi and P. Di Vecchia, Phys. Lett. **400B** (1997) 63, [hep-th/9701190](#).
- [5] M. Frau, I. Pesando, S. Sciuto, A. Lerda and R. Russo, Phys. Lett. **400B** (1997) 52, [hep-th/9702037](#).
- [6] F. Hussain, R. Iengo and C. Núñez, Nucl. Phys. **B497** (1997) 205, [hep-th/9701143](#); F. Hussain, R. Iengo, C. Núñez and C.A. Scrucca, *Closed string radiation from moving D-branes*, [hep-th/9701143](#).
- [7] O. Bergman, M. Gaberdiel and G. Lifschytz, Nucl. Phys. **B509** (1998) 194, [hep-th/9705130](#).
- [8] P. Di Vecchia, M. Frau, I. Pesando, S. Sciuto, A. Lerda, R. Russo, Nucl. Phys. **B507** (1997) 259, [hep-th/9707068](#).
- [9] G. Lifschytz, Phys. Lett. **388B** (1996) 720, [hep-th/9604156](#).

- [10] U. Danielsson, G. Ferretti and I.R. Klebanov, Phys. Rev. Lett. **79** (1997) 1984, hep-th/ 9705084.
- [11] C.P. Bachas, M.R. Douglas and M.B. Green, JHEP **07** (1997) 002, hep-th/ 9705074.
- [12] A. Hanany and E. Witten, Nucl. Phys. **B492** (1997) 152, hep-th/9611230.
- [13] J. Polchinski and A. Strominger, Phys. Lett. **388B** (1996) 736, hep-th/ 9510227.
- [14] S.A. Yost, Nucl. Phys. **B321** (1989) 629.
- [15] M. Bianchi, G. Pradisi and A. Sagnotti, Nucl. Phys. **B376** (1992) 365.
- [16] S. Ramgoolam and L. Thorlacius, Nucl. Phys. **B483** (1997) 248, hep-th/ 9607113.
- [17] C.G. Callan, C. Lovelace, C.R. Nappi and S.A. Yost, Nucl. Phys. **B308** (1988) 221; J. Polchinski and T. Cai, Nucl. Phys. **B286** (1988) 91.
- [18] M.B. Green and Gutperle, Phys. Lett. **398B** (1997) 69, hep-th/9612127.
- [19] T. Kitao, N. Ohta and J.G. Zhou, *Fermionic zero mode and string creation between D4-branes at angles*, hep-th/980135.
- [20] H. Arfaei and M.M. Sheikh-Jabbari, Phys. Lett. **394B** (1997) 288, hep-th/ 9608167; M.M. Sheikh-Jabbari, *Classification of different branes at angles*, hep-th/9710121.
- [21] M. Bertolini, R. Iengo and C.A. Scrucca, *Electric and magnetic interaction of dyonic D-branes and odd spin structure*, hep-th/9801110.
- [22] N. Berkovits, Phys. Rev. Lett. **79** (1997) 1813, hep-th/9706024; Nucl. Phys. **B507** (1997) 731, hep-th/9704109; N. Berkovits and B. Zwiebach, *On the picture dependence of Ramond-Ramond cohomology*, hep-th/9711087.
- [23] V.A. Kostelecky, O. Lechtenfeld, W. Lerche, S. Samuel and S. Watamura, Nucl. Phys **B288** (1987) 173.
- [24] D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. **B271** (1986) 93.